

Publications LCC 1997

Oxidation of pollutants catalyzed by metallophthalocyanines

MEUNIER B., SOROKIN A.

Acc. Chem. Res., 1997, **30**, pp. 470-476 (LCC-4779).

Trigonal planar phosphorus cations

GUERRET O., BERTRAND G.

Acc. Chem. Res., 1997, **30**, pp. 486-493 (LCC-3281).

La chimie pour l'europe et la qualité de la vie

BALAVOINE G.G.A.

Actual. Chim., 1997, pp. 20-21 (LCC-5453).

Multiproperty molecular materials combining conductivity and second-order optical nonlinearities

LACROIX P.G., NAKATANI K.

Adv. Mater., 1997, **9**, pp. 1105-1108 (LCC-4816).

New BETS salts based on magnetic (CuCl_3 , FeCl_4) and non-magnetic (GaCl_4) anions

KUSHCH N.D., DYACHENKO O.A., LYUBOVSKII R.B., PESOTSKII S.I., KARTSOVNIK M.V., KOVALEV A.E., CASSOUX P., KOBAYASHI H.

Adv. Mater. Opt. Electron., 1997, **7**, pp. 57-60 (LCC-4763).

Ruthenium(II) arenes complexes of optically active P.N donor ligands. Synthesis and study of their catalytic activity for the hydrogenation of acetophenone

YANG H., LUGAN N., MATHIEU R.

An. Quim., Int. Ed., 1997, **93**, pp. 28-38 (LCC-4658).

Phosphorus containing dendrimers : synthesis and surface functionalization

MAJORAL J.-P., CAMINADE A.-M., KRAEMER R.

An. Quim., Int. Ed., 1997, **93**, pp. 415-421 (LCC-4808).

Chemoselective polyalkylations of phosphorus-containing dendrimers

LARRE C., CAMINADE A.-M., MAJORAL J.-P.

Angew. Chem. (Int. Ed. Engl.), 1997, **36**, pp. 596-599 (LCC-4592).

Zirconocene-benzyne-mediated intramolecular coupling of bis(alkynyl)phosphane : a way to mono- and tricyclic 1,2-dihydrophosphetes

DUPUIS L., PIRIO N., MEUNIER P., IGAU A., DONNADIEU B., MAJORAL J.-P.

Angew. Chem. (Int. Ed. Engl.), 1997, **36**, pp. 987-989 (LCC-4863).

Generation of alkyl(dicarbonyl)(chloro)-ruthenium dimers in the ruthenium-catalyzed addition of alkyl formates to ethylene

FABRE S., KALCK PH., LAVIGNE G.

Angew. Chem. (Int. Ed. Engl.), 1997, **36**, pp. 1092-1095 (LCC-4852).

Proton-dependent monomer-oligomer interconversion of metal complexes

MATSUMOTO N., MIZUGUCHI Y., MAGO G., EGUCHI S., MIYASAKA H., NAKASHIMA T., TUCHAGUES J.-P.

Angew. Chem. (Int. Ed. Engl.), 1997, **36**, pp. 1860-1862 (LCC-4780).

The first di- μ_3 -hydroxo-bridged tetramanganese(II) complex

STELZIG L., DONNADIEU B., TUCHAGUES J.-P.

Angew. Chem. (Int. Ed. Engl.), 1997, **36**, pp. 2221-2222 (LCC-4809).

Anti-HIV activities of anionic metalloporphyrins and related compounds

SONG R., WITVROUW M., SCHOLS D., ROBERT A., BALZARINI J., DE CLERCQ E., BERNADOU J., MEUNIER B.

Antivir. Chem. Chemother., 1997, **8**, pp. 85-97 (LCC-4593).

Cp₂TiCH₂Si(Me)₂NSiMe₃ : a single-source precursor to titanium-based ceramic thin films by chemical vapor deposition

CHANSOU B., CHOUKROUN R., VALADE L.

Appl. Organomet. Chem., 1997, **11**, pp. 195-203 (LCC-4594).

Nuclease activity and binding characteristics of a cationic "manganese porphyrin-bis(benzimidazole) dye (Hoechst 33258)" conjugate

FRAU S., BERNADOU J., MEUNIER B.

Bioconjugate Chem., 1997, **8**, pp. 222-231 (LCC-4595).

DNA binding and cleavage by a cationic manganese porphyrin-peptide nucleic acid conjugate
BIGEY P.

Bioconjugate Chem., 1997, **8**, pp. 267-270 (LCC-4637).

Organophosphorus compounds : 123. Trimethylsilylphosphaacetylene as cycloaddition partner for 1,3-dipoles

FIEDLER W., REGITZ M., BERTRAND G.

Bull. Soc. Chim. Belg., 1997, **106**, pp. 455-460 (LCC-4319).

One-pot synthesis and reactivity of new tetrazoles, 1,3,4-oxa- and 1,2,3-thiadiazoles

BOURISSOU D., DUPUCH C., DAHAN F., BERTRAND G.

Bull. Soc. Chim. Belg., 1997, **106**, pp. 533-539 (LCC-4546).

Chiral biphosphole iron complexes. Synthesis and characterization of the biphosphaferrocene [Fe(η^5 -C₅H₅)(η^5 -C₄HMe₂P)]₂. X-ray characterization of the dinuclear complex [Fe(η -C₅H₅)(C₄HMe₂PhP)(CO)]₂

DUPUIS A., GOUYGOU M., DARAN J.-C., BALAVOINE G.G.A.

Bull. Soc. Chim. Fr., 1997, **134**, pp. 357-363 (LCC-5454).

Interaction of artemisinin (qinghaosu) with the tetraphenylporphyrinato-manganese(II) complex
ROBERT A., BOULARAN M., MEUNIER B.

C. R. Acad. Sci., Paris, Sér.IIb, 1997, **324**, pp. 59-66 (LCC-4597).

Homogeneous and supported titanates as catalysts for transesterification of acrylic esters

BLANDY C., PELLEGATTA J.-L., CASSOUX P.

Catal. Lett., 1997, **43**, pp. 139-142 (LCC-4628).

Stereoselective synthesis of 2-phosphino-1-zirconaindenes and related 2-phosphinophospholes and stiboles

MIQUEL Y., IGAU A., DONNADIEU B., MAJORAL J.-P., DUPUIS L., PIRIO N., MEUNIER P.

Chem. Commun., 1997, pp. 279-280 (LCC-4675).

Diphosphines with large natural bite angles lead to the formation of non-classical cis-(diphosphine)₂Ru(H)(H²)⁺ complexes

KRANENBURG M., KAMER P.C.J., VAN LEEUWEN P.W.N.M., CHAUDRET B.

Chem. Commun., 1997, pp. 373-374 (LCC-4638).

Homobimetallic vanadium d¹-d¹ complexes : structure and magnetism

CHOUKROUN R., DONNADIEU B., Malfant I., HAUBRICH S., FRANTZ R., GUERIN CH.

Chem. Commun., 1997, pp. 2315-2316 (LCC-4810).

Oxidation of tertiary diaryl catalyzed by a water-soluble metalloporphyrin : C_{aliph}-C_{aliph} versus C_{aliph}-C_{Ar} bond cleavage

WIETZERBIN K., MEUNIER B., BERNADOU J.

Chem. Commun., 1997, pp. 2321-2322 (LCC-4848).

New routes to free and coordinated 1-H-diphosphirenes

BOURISSOU D., CANAC Y., COLLADO M.I., BACEIREDO A., BERTRAND G.

Chem. Commun., 1997, pp. 2399-2400 (LCC-4851).

Zirconaindane phospholanes : key reagents for the synthesis of mono- or tri-cyclic phosphanes and related species

ZABLOCKA M., IGAU A., DONNADIEU B., MAJORAL J.-P., SKOWRONSKA A., MEUNIER P.
Chem. Commun., 1997, pp. 12339-1240 (LCC-4862).

The stable (phosphino)(silyl)carbene as a useful building block: synthesis and reactivity of 2-phosphorus-substituted 2H-azirines

PIQUET V., BACEIREDO A., GORNITZKA H., DAHAN F., BERTRAND G.
Chem. Eur. J., 1997, **3**, pp. 1757-1764 (LCC-4782).

Layered MPS₃ (M=Mn, Cd) thin films as host matrixes for nonlinear optical material processing

LAGADIC I., LACROIX P.G., CLEMENT R.
Chem. Mater., 1997, **9**, pp. 2004-2012 (LCC-4781).

Hydroamination of alkenes. Amidorhodium catalysts

BRUNET J.-J.
Gazz. Chim. Ital., 1997, **127**, pp. 111-118 (LCC-4846).

Spin crossover in the 2,2'-bipyrimidine- (bpym-) bridged iron(II) complexes [Fe(L)(NCX)₂]₂² (bpym) (L=2,2'-bithiazoline (bt) and bpym; X=S, Se). X-ray absorption spectroscopy, magnetic susceptibility, calorimetric, and Mössbauer spectroscopy studies

REAL J.A., CASTRO I., BOUSSEKSOU A., VERDAGUER M., BURRIEL R., CASTRO M., LINARES J., VARRET F.

Inorg. Chem., 1997, **36**, pp. 455-464 (LCC-4625).

Solid state and solution study of trinuclear (Ni, Ba, Ni) complexes : (L¹₂Ni)₂Ba(ClO₄)₂·2H₂O(1) and (L²₂Ni)₂Ba(ClO₄)₂·2H₂O(2) (L¹ = 3-methoxysalicylaldiminato and L² = 3-(2-methoxyethoxy)salicylaldiminato). Crystal and molecular structure of 2

COSTES J.-P., LAURENT J.-P., CHABERT PH., COMMENGES G., DAHAN F.

Inorg. Chem., 1997, **36**, pp. 656-660 (LCC-4600).

Transition-metal derivatives of a functionalized cyclopentadienyl ligand. 15. Synthesis and structures of amino cyclopentadienyl derivatives of rhodium(I) and rhodium(III) including water-soluble compounds

PHILIPPOPOULOS A.I., HADJILIADIS N., HART C.E., DONNADIEU B., MC GOWAN P.C., POILBLANC R.

Inorg. Chem., 1997, **36**, pp. 1842-1849 (LCC-4641).

Versatile complexation ability of very large phosphino-terminated dendrimers

SLANY M., BARDAJI M., CAMINADE A.-M., CHAUDRET B., MAJORAL J.-P.

Inorg. Chem., 1997, **36**, pp. 1939-1945 (LCC-4601).

Spin transition in [Fe(DPEA)(NCS)₂], a compound with the new tetradentate ligand (2-aminoethyl)bis(2-pyridylmethyl)amine (DPEA) : crystal structure, magnetic properties, and Mössbauer spectroscopy

MATOUZENKO G.S., BOUSSEKSOU A., LECOCQ S., VAN KONINSBRUGGEN P.J., PERRIN M., KAHN O., COLLET A.

Inorg. Chem., 1997, **36**, pp. 2975-2981 (LCC-4811).

A general route to strictly dinuclear Cu(II)/Ln(III) complexes. Structural determination and magnetic behavior of two Cu(II)/Gd(III) complexes

COSTES J.-P., DAHAN F., DUPUIS A., LAURENT J.-P.

Inorg. Chem., 1997, **36**, pp. 3429-3433 (LCC-4784).

Origin of the oxygen atom in C-H bond oxidations catalyzed by a water-soluble metalloporphyrin

BALAHURA R.J., SOROKIN A., BERNADOU J., MEUNIER B.

Inorg. Chem., 1997, **36**, pp. 3488-3492 (LCC-4785).

Experimental evidence of a ferromagnetic ground state ($S=9/2$) for a dinuclear Gd(III)-Ni(II) complex

COSTES J.-P., DAHAN F., DUPUIS A., LAURENT J.-P.

Inorg. Chem., 1997, **36**, pp. 4284-4286 (LCC-4786).

Synthesis, crystal structure, magnetic properties, and EPR of $\text{Cu}(\text{bp3ca})\text{Br}_2 \cdot \text{H}_2\text{O}$ (bp3ca=2,2'-bipyridine-3,3'-dicarboxylic acid. Ferromagnetic interactions via unsymmetrical bromide bridges
MENON S., RAJASEKHARAN M.V., TUCHAGUES J.-P.
Inorg. Chem., 1997, **36**, pp. 4341-4346 (LCC-4787).

Magnetic properties of a series of trinuclear complexes $(\text{CuL})_2\text{Mn} \cdot x\text{B}$ (l representing the deprotonated form of *N*- (4-methyl-6-oxo-3-azahept-4-enyl) oxamic acid and B representing respectively H_2O ($x=5,4,5,3,1$), $(\text{CH}_3)_2\text{SO}$ ($x=2$), and $\text{C}_5\text{H}_5\text{N}$ ($x=4$). Crystal and molecular structure of $(\text{CuL})_2\text{Mn} \cdot 2(\text{CH}_3)_2\text{SO}$
COSTES J.-P., LAURENT J.-P., MORENO-SANCHEZ J.M., SUAREZ-VARELA J., AHLGREN M., SUNDBERG M.
Inorg. Chem., 1997, **36**, pp. 4641-4646 (LCC-4788).

Reactivity of the acyclic diazadioxo redox active ligand $[(\text{C}_5\text{H}_5)\text{Fe}(\text{C}_5\text{H}_4\text{CH}_2\text{N}(\text{CH}_3)(\text{CH}_2)_2\text{OCH}_2)_2]$: NMR, electrochemical, and Mössbauer studies. Crystal structure of its copper complex
DELAVAUX-NICOT B., BIGEARD A., BOUSSEKSOU A., DONNADIEU B., COMMENGES G.
Inorg. Chem., 1997, **36**, pp. 4789-4797 (LCC-4789).

Synthesis and characterization of $[\text{acridinium}][\text{Ni}(\text{dmit})_2]_3$ and $[\text{phenazinium}][\text{Ni}(\text{dmit})_2]_3$ (dmit=2-thioxo-1,3-dithiole-4,5-dithiolate)
VELDHUIZEN Y.S.J., SMEETS W.J.J., VELDMAN N., SPEK A.L., FAULMANN C., AUBANSENZIER P., JEROME D., PAULUS P.M., HAASNOOT J.G., REEDIJK J.
Inorg. Chem., 1997, **36**, pp. 4930-4937 (LCC-4790).

Polymorphism in spin transition systems. Crystal structure, magnetic properties, and Mössbauer spectroscopy of three polymorphic modifications of $[\text{Fe}(\text{DPPA})(\text{NCS})_2]$ [DPPA = (3-aminopropyl)bis(2-pyridylmethyl)amine]
MATOUZENKO G.S., BOUSSEKSOU A., LECOCQ S., VAN KONINGSBRUGGEN P.J., PERRIN M., KAHN O., COLLET A.
Inorg. Chem., 1997, **36**, pp. 5869-5879 (LCC-4812).

Synthesis, structure, and magnetic and redox properties of linear bis-dinuclear complexes afforded by Schiff base ligands containing catecholate and pyridine or imidazole groups
THEIL S., YERANDE R., CHIKATE R., DAHAN F., BOUSSEKSOU A., PADHYE S., TUCHAGUES J.-P.
Inorg. Chem., 1997, **36**, pp. 6279-6286 (LCC-4813).

Orientation-selective ESEEM study and crystal structure of a $\text{Cu}^{\text{I}}(\text{thiochrome})\text{Cl}_2$ complex
LOULUDI M., DELIGIANNAKIS Y., TUCHAGUES J.-P., DONNADIEU B., HADJILIADIS N.
Inorg. Chem., 1997, **36**, pp. 6335-6342 (LCC-4814).

Modulation of quantum mechanical exchange couplings in transition metal hydrides through hydrogen bonding
AYLLON J.A., SABO-ETIENNE S., CHAUDRET B., ULRICH S., LIMBACH H.-H.
Inorg. Chim. Acta, 1997, **259**, pp. 1-4 (LCC-4895).

Nitrilimines: evidence for the allenic structure in solution, experimental and ab initio studies of the barrier to racemization, and first diastereoselective [3+2]-cycloaddition
FAURE J.-L., REAU R., WONG M.W., KOCH R., WENTRUP C., BERTRAND G.
J. Am. Chem. Soc., 1997, **119**, pp. 2819-2824 (LCC-4669).

Identification of Zr(III) allyl species in the Negishi system $[\text{Cp}_2\text{ZrCl}_2, 2\text{LiBu}]$. An ESR study
SOLEIL F., CHOUKROUN R.
J. Am. Chem. Soc., 1997, **119**, pp. 2938-2939 (LCC-4602).

Synthesis and characterization of chelating bis(silane) complexes $[\text{RuH}_2\{(\eta^2\text{-HSiMe}_2)_2\text{X}\}(\text{PCy}_3)_2]$ ($\text{X}=\text{C}_6\text{H}_4, \text{O}$) containing two Ru-(η^2 -Si-H) bonds
DELPECH F., SABO-ETIENNE S., CHAUDRET B., DARAN J.-C.
J. Am. Chem. Soc., 1997, **119**, pp. 3167-3168 (LCC-4603).

- Reversible migratory insertion/ β -alkyl elimination in α -agostic alkylniobium alkyne complexes
ETIENNE M., MATHIEU R., DONNADIEU B.
J. Am. Chem. Soc., 1997, **119**, pp. 3218-3228 (LCC-4610).
- Characterization of the first covalent adduct between artemisinin and a heme model
ROBERT A., MEUNIER B.
J. Am. Chem. Soc., 1997, **119**, pp. 5968-5969 (LCC-4791).
- 1,2,4-triazole-3,5-diylidene : a building block for organometallic polymer synthesis
GUERRET O., SOLE S., GORNITZKA H., TEICHERT M., TRINQUIER G., BERTRAND G.
J. Am. Chem. Soc., 1997, **119**, pp. 6668-6669 (LCC-4351).
- A room temperature stable 1,3-diphosphallyl radical
CANAC Y., BACEIREDO A., SCHOELLER W., GIGMES D., BERTRAND G.
J. Am. Chem. Soc., 1997, **119**, pp. 7579-7580 (LCC-4792).
- Synthesis, structural, magnetic, and redox properties of asymmetric diiron complexes with a single terminally bound phenolate ligand. Relevance to the purple acid phosphatase enzymes
LAMBERT E., CHABUT B., CHARDON-NOBLAT S., DERONZIER A., CHOTTARD G., BOUSSEKSOU A., TUCHAGUES J.-P., LAUGIER J., BARDET M., LATOUR J.-M.
J. Am. Chem. Soc., 1997, **119**, pp. 9424-9437 (LCC-4793).
- 1,2,4,5-tetra- σ -diphosphetes ($x=2-6$) : on the persistence of the P-P bond
SANCHEZ M., REAU R., GORNITZKA H., DAHAN F., REGITZ M., BERTRAND G.
J. Am. Chem. Soc., 1997, **119**, pp. 9720-9728 (LCC-3204).
- Synthesis and reactivity of a stable n^2 - (diphosphirenylium) $W(CO)_5$ complex
BOURISSOU D., CANAC Y., COLLADO M.I., BACEIREDO A., BERTRAND G.
J. Am. Chem. Soc., 1997, **119**, pp. 9923-9924 (LCC-4794).
- Superconductor-to-insulator transition in an organic metal incorporating magnetic anions : 1-(BETS) $_2$ (Fe $_x$ Ga $_{1-x}$)Cl $_4$ [BETS=bis(ethylenedithio)tetraselenafulvalene ; $x=0,55$ and $0,43$
KOBAYASHI H., SATO A., ARAI E., AKUTSU H., KOBAYASHI A., CASSOUX P.
J. Am. Chem. Soc., 1997, **119**, pp. 12392-12393 (LCC-4820).
- Highly correlated organic conductor with magnetic anions exhibiting a p-d coupled metal-insulator transition, 1-(BETS) $_2$ FeBr $_x$ Cl $_{4-x}$ (BETS=bis(ethylenedithio)tetraselenafulvalene)
AKUTSU H., ARAI E., KOBAYASHI H., TANAKA H., KOBAYASHI A., CASSOUX P.
J. Am. Chem. Soc., 1997, **119**, pp. 12681-12682 (LCC-4821).
- Synthesis of water-soluble ruthenium porphyrins as DNA cleavers and potential cytotoxic agents
HARTMANN M., ROBERT A., DUARTE V., KEPPLER B.K., MEUNIER B.
J. Biol. Inorg. Chem., 1997, **2**, pp. 427-432 (LCC-4795).
- Synthesis, crystal structure, conductivity and magnetic properties of trifluoromethylated dinuclear copper(II) complexes with tetracyanoquinodimethane
LACROIX P.G., DARAN J.-C.
J. Chem. Soc., Dalton Trans., 1997, pp. 1369-1374 (LCC-4627).
- Exchange couplings and hydrogen dynamics in ruthenium trihydride adducts with coinage-metal cations
MANZANO B., JALON F., MATTHES J., SABO-ETIENNE S., CHAUDRET B., ULRICH S., LIMBACH H.-H.
J. Chem. Soc., Dalton Trans., 1997, pp. 3153-3159 (LCC-4891).
- Oxidative damage on RNA mediated by cationic metalloporphyrin-antisense oligonucleotides conjugates
DUARTE V., SIXOU S., FAVRE G., PRATVIEL G., MEUNIER B.
J. Chem. Soc., Dalton Trans., 1997, pp. 4113-4118 (LCC-4796).

- Studies of the interrelated series of triiridatrithia-cyclohexane and trimethylatrithia (or trioxo) - adamantane rhodium and iridium complexes
DEVILLERS J., MAISONNAT A., POILBLANC R.
J. Clust. Sci., 1997, **8**, pp. 1-26 (LCC-4642).
- Metallophthalocyanine-catalyzed oxidation of catechols by H₂O₂ and its surrogates
SOROKIN A., FRAISSE L., RABION A., MEUNIER B.
J. Mol. Catal. A, 1997, **117**, pp. 103-114 (LCC-4611).
- N*-phosphino- and *N*-phosphinonitrilimines: from nucleophilic to electrophilic 1,3-dipoles
PALACIOS F., PAGALDAY J., PIQUET V., DAHAN F., BACEIREDO A., BERTRAND G.
J. Org. Chem., 1997, **62**, pp. 292-296 (LCC-4662).
- Metalloporphyrin-catalyzed oxidation of 2-methylnaphthalene to vitamin K₃ and 6-methyl-1,4-naphthoquinone by potassium monopersulfate in aqueous solution
SONG R., SOROKIN A., BERNADOU J., MEUNIER B.
J. Org. Chem., 1997, **62**, pp. 673-678 (LCC-4612).
- Dual radical/polar Pudovik reaction : application field of new activation methods
SEMENZIN D., ETEMAD-MOGHADAM G., ALBOUY D., DIALLO O.S., KOENIG M.
J. Org. Chem., 1997, **62**, pp. 2414-2422 (LCC-5500).
- Preparation of tetracationic metalloporphyrin-spermine conjugates
JAKOBS A., BERNADOU J., MEUNIER B.
J. Org. Chem., 1997, **62**, pp. 3505-3510 (LCC-4753).
- Phosphate-, phosphite-, ylido-, and phosphonate-terminated dendrimers
PREVOTE D., CAMINADE A.-M., MAJORAL J.-P.
J. Org. Chem., 1997, **62**, pp. 4834-4841 (LCC-4797).
- Facile synthesis of amino-functionalised cyclopentadienyl sodium compounds and subsequent formation of rhodium complexes : proton-deuterium exchange for the sodium salts
MC GOWAN P.C., HART C.E., DONNADIEU B., POILBLANC R.
J. Organomet. Chem., 1997, **528**, pp. 191-194 (LCC-4643).
- Synthesis of bowl-shaped dendrimers from generation 1 to generation 8
LAUNAY N., CAMINADE A.-M., MAJORAL J.-P.
J. Organomet. Chem., 1997, **529**, pp. 51-58 (LCC-4613).
- The versatile behavior of a 1,2,1⁵-azaphosphete towards protic nucleophiles
BOUHADIR G., BIEGER K., LIVOTTO P., REAU R., GORNITZKA H., DAHAN F., BERTRAND G.
J. Organomet. Chem., 1997, **529**, pp. 79-85 (LCC-4754).
- Transition-metal complexes of cyclopentadienylphosphines XIV. Reduction reactions of various derivatives of the chelate fragment $[\{h^5:h^1-C_5H_4(CH_2)_2PPh_2\}M^{III}]^{2+}$, M=Rh, Ir
LEE I., DAHAN F., MAISONNAT A., POILBLANC R.
J. Organomet. Chem., 1997, **532**, pp. 159-170 (LCC-4644).
- Tandem acylation-complexation of a chlorophosphine by carbonylferrates
BRUNET J.-J., CAPPERUCCI A., CHAUVIN R., DONNADIEU B.
J. Organomet. Chem., 1997, **533**, pp. 79-81 (LCC-4777).
- A rhodium(I) amide dimer : synthesis, structural characterisation and preliminary reactivity studies
BRUNET J.-J., DARAN J.-C., NEIBECKER D., ROSENBERG L.
J. Organomet. Chem., 1997, **538**, pp. 251-256 (LCC-4847).
- Experimental evidence of the symmetry breakdown in 4,8-diethoxy-2,7-bis(benzo-1',3'-dithiol-2'-ylidene)-1,3,5,7-tetrathia-*s*-indacene compounds
POKHODNIA K., CASSOUX P., BONVOISIN J., MLAYAH A., BROSSARD L., FRENZEL S., MULLEN K.
J. Phys. Chem., 1997, **101**, pp. 3665-3670 (LCC-4674).

Surface spectroscopic study of carbon monoxide adsorption on nanoscale nickel colloids prepared from a zerovalent organometallic precursor

DE CARO D., BRADLEY J.S.

Langmuir, 1997, **13**, pp. 3067-3069 (LCC-5486).

L'anion $[\text{HFe}(\text{CO})_4]^-$, précurseur pour une chimie de coordination originale

BRUNET J.-J., CHAUVIN R., NEIBECKER D.

Lett. Sci. Chim., 1997, **59**, pp. 8-10 (LCC-5304).

Large dipole moments of phosphorus-containing dendrimers

LARTIGUE M.-L., DONNADIEU B., GALLIOT C., CAMINADE A.-M., MAJORAL J.-P.

Macromolecules, 1997, **30**, pp. 7335-7337 (LCC-4854).

Progress in research and development for high temperature and low temperature superconductors

BARDAJI M., SLANY M., LARTIGUE M.-L., CAMINADE A.-M., CHAUDRET B., MAJORAL J.-P.

Main Group Chem., 1997, **2**, pp. 133-140 (LCC-3149).

Inhibition of in vitro translation by antisense oligonucleotides covalently linked to a nucleic acid cleaver based on a cationic manganese porphyrin motif, Mn-trisMPyP

DUARTE V., PRATVIEL G., MEUNIER B., BERTON M., SIXOU S., FAVRE G.

New J. Chem., 1997, **21**, pp. 55-60 (LCC-4614).

Three-dimensional bimaocyclic phosphorus compounds : phosphorus cryptands

CAMINADE A.-M., KRAEMER R., MAJORAL J.-P.

New J. Chem., 1997, **21**, pp. 627-632 (LCC-4623).

Synthesis, characterization, and some chemistry of $(\text{bp}^{\text{F}})\text{RuH}(\text{COD})$ ($\text{bp}^{\text{F}} = \text{H}_2\text{B}[\text{N}_2\text{C}_3\text{H}(\text{CF}_3)_2]$), an electrophilic hydrido ruthenium complex

RODRIGUEZ V., FULL J., DONNADIEU B., SABO-ETIENNE S., CHAUDRET B.

New J. Chem., 1997, **21**, pp. 847-849 (LCC-4893).

Solid state and solution studies of the compounds resulting from the equimolecular condensation of a diamine with a dione monoxime

COSTES J.-P., DAHAN F., DUPUIS A., LAURENT J.-P.

New J. Chem., 1997, **21**, pp. 1211-1217 (LCC-4798).

Synthesis of platinum nanoparticles stabilized by CO and tetrahydrothiophene. Facile conversion to molecular species

BARDAJI M., VIDONI O., RODRIGUEZ A., AMIENS C., CHAUDRET B., CASANOVE M.-J., LECANTE P.

New J. Chem., 1997, **21**, pp. 1243-1249 (LCC-4892).

Catalysis of peroxyxynitrite reactions by manganese and iron porphyrins

BALAVOINE G.G.A., GELETII Y.V., BEJAN D.

Nitric Oxide, 1997, **1**, pp. 507-521 (LCC-3215).

Influence of the nature of the porphyrin ligand on the nuclease activity of metalloporphyrin-oligonucleotide conjugates designed with cationic, hydrophobic or anionic metalloporphyrins

MESTRE B., PITIE M., LOUP C., CLAPAROLS C., PRATVIEL G., MEUNIER B.

Nucleic Acids Res., 1997, **25**, pp. 1022-1027 (LCC-4615).

Iron-gold (or-mercury) carbide clusters derived from $[\text{Fe}_6\text{C}(\text{CO})_{16}]^{2-}$. X-ray crystal structures of $(\text{NEt}_4)[\text{Fe}_6\{\text{AuPPh}_3\}(\text{CO})_{16}]$ and $[\text{Fe}_4\text{C}\{\text{AUPPh}_3\}(\text{CO})_{11}(\text{NO})]$

ROSSELL O., SECO M., SEGALES G., ALVAREZ S., PELLINGHELLI M.-A., TIRIPICCHIO A., DE MONTAUZON D.

Organometallics, 1997, **16**, pp. 236-245 (LCC-4616).

Phosphorus-containing dendrimers as multidentate ligands : palladium, platinum, and rhodium complexes

BARDAJI M., KUSTOS M., CAMINADE A.-M., MAJORAL J.-P., CHAUDRET B.

Organometallics, 1997, **16**, pp. 403-410 (LCC-4617).

Biphosphole : a C_2 symmetry chiral bidentate ligand. Synthesis and characterization of nickel, palladium, platinum, and rhodium complexes

GOUYGOU M., TISSOT O., DARAN J.-C., BALAVOINE G.G.A.

Organometallics, 1997, **16**, pp. 1008-1015 (LCC-4622).

Ruthenium(II) complexes containing optically active hemilabile P,N,O-tridentate ligands. Synthesis and evaluation in catalytic asymmetric transfer hydrogenation of acetophenone by propan-2-ol

YANG H., ALVAREZ-GRESSIER M., LUGAN N., MATHIEU R.

Organometallics, 1997, **16**, pp. 1401-1409 (LCC-4618).

First NMR observation of the intermolecular dynamic proton transfer equilibrium between a hydride and coordinated dihydrogen : $(dppm)_2HRuH...H-OR=[(dppm)_2HRu(H_2)]^+(OR)^-$

AYLLON J.A., GERVAUX C., SABO-ETIENNE S., CHAUDRET B.

Organometallics, 1997, **16**, pp. 2000-2002 (LCC-4639).

Study of the bonding properties of the new ligands $C_5H_3N(2-R')$ (6- CH_2 PPhR) toward rhodium(I). Evidence for a dynamic competition for bonding between O and N-donor centers when $R=O$ -anisyl, $R'=Me$

LUGAN N., MATHIEU R., YANG H.

Organometallics, 1997, **16**, pp. 2089-2095 (LCC-4619).

Versatile behavior of a new class of bicyclic compounds : zirconacyclopentadiene phosphiranes

MAHIEU A., MIQUEL Y., IGAU A., DONNADIEU B., MAJORAL J.-P.

Organometallics, 1997, **16**, pp. 3086-3088 (LCC-4860).

Ruthenium hydride and dihydrogen complexes with dendrimeric multidentate ligands

BARDAJI M., CAMINADE A.-M., MAJORAL J.-P., CHAUDRET B.

Organometallics, 1997, **16**, pp. 3489-3497 (LCC-4799).

A novel route to 2-cyclohexenones via reaction of the manganese carbene anions $[(\eta^5-MeC_5H_4)(CO)_2Mn=C(OEt)CHR^-]$ ($R=H, Me$) with α, β -unsaturated ketones

MONGIN C., LUGAN N., MATHIEU R.

Organometallics, 1997, **16**, pp. 3873-3875 (LCC-4911).

[Cp₂M] metallocene species promoted reductive coupling of $Cl_2C=PR$: synthesis, structure, and reactivity of the first 3,4-dihalo-1,2-dihydro-1,2-diphosphetes

CENAC N., CHROSTOWSKA A., SOTIROPOULOS J.-M., DONNADIEU B., IGAU A., PFISTER-GUILLOUZO G., MAJORAL J.-P.

Organometallics, 1997, **16**, pp. 4551-4556 (LCC-4800).

Disproportionation of cationic zirconium complexes : a possible pathway to the deactivation of catalytic cationic systems

CHOUKROUN R., DOUZIECH B., DONNADIEU B.

Organometallics, 1997, **16**, pp. 5517-5521 (LCC-5508).

Synthesis and reactivity of dendrimers based on phosphoryl (P=O) groups

LARTIGUE M.-L., CAMINADE A.-M., MAJORAL J.-P.

Phosph. Sulf. Silicon, 1997, **123**, pp. 21-34 (LCC-4909).

Synthesis and characterization of manganese(III) complexes of a chiral disulfonamide ligand based on trans-1,2-diaminocyclohexane

BERNARDO K., ROBERT A., MEUNIER B.

Polyhedron, 1997, **16**, pp. 2365-2368 (LCC-4648).

Regioselective stepwise growth of dendrimer units in the internal voids of a main dendrimer

GALLIOT C., LARRE C., CAMINADE A.-M., MAJORAL J.-P.

Science, 1997, **277**, pp. 1981-1984 (LCC-4801).

A stable molecular metal with a binuclear magnetic anion, α -(BETS)₄Cu₂Cl₆
KOBAYASHI A., SATO A., ARAI E., KOBAYASHI H., FAULMANN C., KUSHCH N.D., CASSOUX P.

Solid State Commun., 1997, **103**, pp. 371-374 (LCC-4802).

Easy preparation of Collman-type reagents: reaction of [HFe(CO)₄]⁻ with tris(dimethylamino)phosphine
BRUNET J.-J., CHAUVIN R., NEIBECKER D.

Synth. Commun., 1997, **27**, pp. 1433-1437 (LCC-4778).

Structural and electrical properties of BETS compounds with transition metal complex anions
KOBAYASHI H., ARAI E., SATO A., NAITO T., TANAKA H., KOBAYASHI A., SAITO T., CASSOUX P.

Synth. Met., 1997, **85**, pp. 1595-1596 (LCC-4630).

Interplay of conductivity, magnetism (and nonlinear optics) in organic-inorganic compounds
BROSSARD L., CASSOUX P., DYACHENKO O.A., FAULMANN C., GRAJA A., KOBAYASHI H., KOBAYASHI A., KUSHCH N.D., LACROIX P.G., LEGROS J.-P., MALFANT I., POKHODNIA K., TOKUMOTO M.

Synth. Met., 1997, **86**, pp. 1785-1788 (LCC-4760).

Spectroscopic investigation of the symmetry breakdown in bis-TTF compounds
POKHODNIA K., CASSOUX P., BONVOISIN J., MLAYAH A., BROSSARD L., FRENZEL S., MULLEN K.

Synth. Met., 1997, **86**, pp. 1999-2000 (LCC-4761).

Microreflectance infrared study of the organic conductor κ -(BETS)₂FeCl₄
OLEJNICZAK I., GRAJA A., KUSHCH N.D., CASSOUX P., KOBAYASHI H.

Synth. Met., 1997, **86**, pp. 2155-2156 (LCC-4762).

Magnetic anisotropy of organic conductor λ -(BETS)₂FeCl₄
TOKUMOTO M., NAITO T., KOBAYASHI H., KOBAYASHI A., LAUKIN V.N., BROSSARD L., CASSOUX P.

Synth. Met., 1997, **86**, pp. 2161-2162 (LCC-4050).

Application of the Horner-Wadsworth-Emmons reaction to the functionalization of dendrimers: synthesis of amino acid terminated dendrimers

PREVOTE D., LE ROY-GOURVENNEC S., CAMINADE A.-M., MASSON S., MAJORAL J.-P.

Synthesis, 1997, pp. 1199-1207 (LCC-4803).

Chiroptical properties of dendrimers with stereogenic end groups

LARTIGUE M.-L., CAMINADE A.-M., MAJORAL J.-P.

Tetrahedron Asymmetry, 1997, **8**, pp. 2697-2708 (LCC-4804).

New homochiral bis(oxazoline) ligands for asymmetric catalysis

HOARAU O., AIT-HADDOU H., CASTRO M., BALAVOINE G.G.A.

Tetrahedron Asymmetry, 1997, **8**, pp. 3755-3764 (LCC-4855).

A new method for the regio and stereoselective hydrocarboxylation of alkynes using NaHFe(CO)₄/CH₂Cl₂ system

PERIASAMY M., RADHAKRISHNAN U., RAMESHKUMAR C., BRUNET J.-J.

Tetrahedron Lett., 1997, **38**, pp. 1623-1626 (LCC-2087).

[Cp₂ZrHCl]_n a useful reducing agent in phosphorus chemistry

ZABLOCKA M., DELEST B., IGAU A., SKOWRONSKA A., MAJORAL J.-P.

Tetrahedron Lett., 1997, **38**, pp. 5997-6000 (LCC-4861).