

## **Publications LCC 1998**

Key intermediates in the synthesis of 2,2'-biphosphole

TISSOT O., GOUYGOU M., DARAN J.-C., BALAVOINE G.G.A.  
*Acta Crystallogr., Sect. C*, 1998, **54**, pp. 676-679 (LCC-4857).

A new Ni(dmit)<sub>2</sub> salt with a paramagnetic cation : [Cu(bpy)<sub>3</sub>][Ni(C<sub>3</sub>S<sub>5</sub>)<sub>2</sub>]<sub>2</sub>

FAULMANN C., VELDHUIZEN Y.S.J., HAASNOOT J.G., REEDIJK J., CASSOUX P.  
*Acta Crystallogr., Sect. C*, 1998, **54**, pp. 1827-1830 (LCC-5199).

Ylidic four p electron four-membered 1<sup>5</sup>-phosphorus heterocycles : electronical isomers of heterocyclobutadienes

BERTRAND G.  
*Angew. Chem. (Int. Ed. Engl.)*, 1998, **37**, pp. 270-281 (LCC-4823).

The azide-nitrilimine analogy in aluminum chemistry

EMIG N., GABBAI F.P., KRAUTSCHEID H., REAU R., BERTRAND G.  
*Angew. Chem. (Int. Ed. Engl.)*, 1998, **37**, pp. 989-992 (LCC-5158).

Highly enantioselective one-pot synthesis of chiral tri- and tetrasubstituted ferrocenes from 1,1'-ferrocenedicarbaldehyde

IFTIME G., DARAN J.-C., MANOURY E., BALAVOINE G.G.A.  
*Angew. Chem. (Int. Ed. Engl.)*, 1998, **37**, pp. 1698-1701 (LCC-5461).

Calixarene-based copper(I) complexes as models for monocopper sites in enzymes

BLANCHARD S., LE CLAINCHE L., RAGER M.-N., CHANSOU B., TUCHAGUES J.-P., DUPRAT A.F., LE MEST Y., REINAUD O.  
*Angew. Chem. (Int. Ed. Engl.)*, 1998, **37**, pp. 2732-2735 (LCC-5200).

Cationic homoleptic vanadium(II), (IV), and (V) complexes arising from protonolysis of [V(NEt<sub>2</sub>)<sub>4</sub>]

CHOUKROUN R., MOUMBOKO P., CHEVALIER S., ETIENNE M., DONNADIEU B.  
*Angew. Chem. (Int. Ed. Engl.)*, 1998, **37**, pp. 3169-3172 (LCC-5201).

Précurseurs métallorganiques pour le dépôt chimique à partir d'une phase gazeuse de revêtements dans le système Ti-V-C-N

MAURY F., TEYSSANDIER F., VALADE L.  
*Ann. Chim.-Sci. Mater.*, 1998, **23**, pp. 637-653 (LCC-5075).

Problèmes spécifiques liés à la vaporisation des précurseurs organométalliques solides utilisés pour les dépôts dans le système V-C-N

TEYSSANDIER F., POIRIER L., SLIFIRSKI J., VALADE L., DANJOY C., REYNES A., JAUBERTEAU J.-L., SIBIEUDE F.  
*Ann. Chim.-Sci. Mater.*, 1998, **23**, pp. 655-666 (LCC-5076).

Mécanisme de décomposition des composés M(NEt<sub>2</sub>)<sub>4</sub> (M=V, Cr) lors de leur utilisation comme mono-source pour la croissance de films M-C-N par MOCVD

BONNEFOND P., FEURER R., MAURY F., REYNES A., CHANSOU B., CHOUKROUN R., CASSOUX P.  
*Ann. Chim.-Sci. Mater.*, 1998, **23**, pp. 667-679 (LCC-5077).

Possibilités d'obtention de dépôts dans le système céramique Ti-V-C-N à partir de sources moléculaires

VALADE L., CHOUKROUN R., DANJOY C., CHANSOU B., DE CARO D., CASSOUX P.  
*Ann. Chim.-Sci. Mater.*, 1998, **23**, pp. 721-732 (LCC-5078).

Evaluation of the simultaneous use of Cp<sub>2</sub>VMe<sub>2</sub> and CpTiCl<sub>2</sub>N(SiMe<sub>3</sub>)<sub>2</sub> as precursors to ceramic thin films containing titanium and vanadium : towards titanium-vanadium carbinitride

VALADE L., DANJOY C., CHANSOU B., RIVIERE E., PELLEGATTA J.-L., CHOUKROUN R., CASSOUX P.  
*Appl. Organomet. Chem.*, 1998, **12**, pp. 173-187 (LCC-4824).

Coherent and incoherent dihydrogen dynamics in a ruthenium trihydride complex with the tris(pyrollyl)phosphine ligand  
GRUNDEMANN S., LIMBACH H.-H., RODRIGUEZ V., DONNADIEU B., SABO-ETIENNE S., CHAUDRET B.

*Ber. Bunsenges. Physik. Chem.*, 1998, **102**, pp. 344-353 (LCC-4896).

Preparation of a spermine conjugate of the bis-phenanthroline ligand Clip-Phen and evaluation of the corresponding copper complex  
PITIE M., MEUNIER B.

*Bioconjugate Chem.*, 1998, **9**, pp. 604-611 (LCC-5079).

Modification of the thiourea linkage of a fluorescein-oligonucleotide conjugate to a guanidinium motif during ammonia deprotection  
DUBEY I., PRATVIEL G., MEUNIER B.

*Bioconjugate Chem.*, 1998, **9**, pp. 627-632 (LCC-5080).

Synthesis, structure, and properties of di- $\mu$ -phenoxo-bridged dinuclear  $\text{Cu}^{\text{II}}\text{Ni}^{\text{II}}$  and  $\text{Cu}^{\text{II}}\text{Co}^{\text{II}}$  complexes and di- $\mu$ -phenoxo- $\mu$ -amido-bridged tetranuclear  $\text{Cu}_2^{\text{II}}\text{Zn}_2^{\text{II}}$  and  $\text{Cu}_4^{\text{II}}$  complexes derived from the 1,2-bis(2-oxidobenzamidato)benzenecuprate(II) dianionic ligand-complex  
SUNATSUKI Y., MATSUO T., NAKAMURA M., KAI F., MATSUMOTO N., TUCHAGUES J.-P.

*Bull. Chem. Soc. Jpn.*, 1998, **71**, pp. 2611-2619 (LCC-5202).

Phosphorus containing dendrimers : chemistry on the surface and within the cascade structure  
MAJORAL J.-P., CAMINADE A.-M., LAURENT R.

*Bull. Pol. Acad. Sci. -Chem.*, 1998, **46**, pp. 319-329 (LCC-5203).

First "layer-block" dendrimer built with a regular alternation of two types of repeat units up to the fourth generation  
LARTIGUE M.-L., LAUNAY N., DONNADIEU B., CAMINADE A.-M., MAJORAL J.-P.

*Bull. Soc. Chim. Fr.*, 1998, **134**, pp. 981-988 (LCC-4822).

Oxidation of 2,4,6-trichlorophenol (TCP) catalyzed by iron tetrasulfophthalocyanine (FePcS) supported on a cationic ion-exchange resin  
HADASCH A., SOROKIN A., RABION A., FRAISSE L., MEUNIER B.

*Bull. Soc. Chim. Fr.*, 1998, **134**, pp. 1025-1032 (LCC-4828).

Oxidation-induced ring expansion of a 2-phosphino-2H-azirine  
PIQUET V., BACEIREDO A., DAHAN F., BERTRAND G.

*C. R. Acad. Sci. Paris, Série IIc*, 1998, **1**, pp. 123-128 (LCC-5204).

Oxidation of guanine bases mediated by a manganese-porphyrin-oligonucleotide conjugate onto a G-rich DNA target : 8-oxo-7,8-dihydroguanine is not the major lesion  
MESTRE B., NASCIMBEN S., PRATVIEL G., MEUNIER B.

*C. R. Acad. Sci. Paris, Série IIc*, 1998, **1**, pp. 725-736 (LCC-5205).

Oxidation-induced ring expansion of a 2-phosphino-2H-azirine  
PIQUET V., BACEIREDO A., DAHAN F., BERTRAND G.

*C. R. Acad. Sci., Paris, Sér. II c*, 1998, **1**, pp. 123-128 (LCC-4819).

Preparation of cationic non-metallated or zinc-porphyrin-oligonucleotide fluorescent conjugates  
DUBEY I., PRATVIEL G., MEUNIER B.

*C. R. Acad. Sci., Paris, Sér. II c*, 1998, **1**, pp. 259-267 (LCC-4883).

Une famille originale de complexes hétérodinucléaires  $\text{Co(II)-Ln(III)}$  : synthèse et étude magnétostructurale  
COSTES J.-P., DAHAN F., DUPUIS A., LAURENT J.-P.

*C. R. Acad. Sci., Paris, Sér. II c*, 1998, **1**, pp. 417-420 (LCC-5110).

- Versatile pentadentate 1,5-bis(salicylideneamino)pentan-3-ol type ligands yield novel tri- and tetra-manganese(II) complexes : structure and properties  
 STELZIG L., STEINER A., CHANSOU B., TUCHAGUES J.-P.  
*Chem. Commun.*, 1998, pp. 771-772 (LCC-4825).
- One pot synthesis of mono and bicyclic phosphiranes and phosphirenes  
 ZABLOCKA M., MIQUEL Y., IGAU A., MAJORAL J.-P., SKOWRONSKA A.  
*Chem. Commun.*, 1998, pp. 1177-1178 (LCC-5129).
- Design, synthesis and cleaving activity of an abiotic nuclease based on a manganese (III) porphyrin complex bearing two acridine moieties  
 DREXTER C., HOSSEINI M.W., PRATVIEL G., MEUNIER B.  
*Chem. Commun.*, 1998, pp. 1343-1344 (LCC-5168).
- Isolation and characterization of an oxidative degradation product of a polypyridine ligand  
 RENZ M., HEMMERT C., DONNADIEU B., MEUNIER B.  
*Chem. Commun.*, 1998, pp. 1635-1636 (LCC-5062).
- Observing and modelling energetically close  $\alpha$ - and  $\beta$ -carbon-hydrogen agostic interactions in an isopropyl tris(pyrazolyl)boratoniobium complex  
 JAFFART J., MATHIEU R., ETIENNE M., MCGRADY J.E., EISENSTEIN O., MASERAS F.  
*Chem. Commun.*, 1998, pp. 2011-2012 (LCC-5081).
- "Oxo-hydroxo tautomerism" as useful mechanistic tool in oxygenation reactions catalysed by water-soluble metalloporphyrins  
 BERNADOU J., MEUNIER B.  
*Chem. Commun.*, 1998, pp. 2167-2173 (LCC-5325).
- Dramatic increase of the DNA cleavage activity of Cu(Clip-phen) by fixing the bridging linker on the C<sup>3</sup> position of the phenanthroline units  
 PITIE M., SUDRES B., MEUNIER B.  
*Chem. Commun.*, 1998, pp. 2597-2598 (LCC-5206).
- [Cp<sub>2</sub>V( $\mu$ -h<sup>2</sup>:h<sup>4</sup>-butadiyne)MCp'<sub>2</sub>] (M=Ti, Zr; Cp'=C<sub>5</sub>H<sub>5</sub>, C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>): heterodimetallic complexes from a bis(alkynyl)metallocene and vanadocene. Unexpected structure of [Cp<sub>2</sub>V( $\mu$ -h<sup>2</sup>:h<sup>4</sup>-butadiyne)Zr(C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>)<sub>2</sub>] containing two planar-tetracoordinate carbon atoms  
 DANJOY C., ZHAO J.S., DONNADIEU B., LEGROS J.-P., VALADE L., CHOUKROUN R., ZWICK A., CASSOUX P.  
*Chem. Eur. J.*, 1998, **4**, pp. 1100-1105 (LCC-5063).
- Alkylating properties of antimalaric artemisinin derivatives and synthetic trioxanes when activated by a reduced heme model  
 ROBERT A., MEUNIER B.  
*Chem. Eur. J.*, 1998, **4**, pp. 1287-1296 (LCC-5167).
- Nature of the magnetic interaction in the (Cu<sup>2+</sup>, Ln<sup>3+</sup>) pairs : an empirical approach based on the comparison between homologous (Cu<sup>2+</sup>, Ln<sup>3+</sup>) and (Ni<sup>2+</sup><sub>LS</sub>, Ln<sup>3+</sup>) complexes  
 COSTES J.-P., DAHAN F., DUPUIS A., LAURENT J.-P.  
*Chem. Eur. J.*, 1998, **4**, pp. 1616-1620 (LCC-5082).
- Regioselective gold complexation within the cascade structure of phosphorus-containing dendrimers  
 LARRE C., DONNADIEU B., CAMINADE A.-M., MAJORAL J.-P.  
*Chem. Eur. J.*, 1998, **4**, pp. 2031-2036 (LCC-5085).
- Temperature-composition phase diagram of the organic alloys, 1-BETS<sub>2</sub>(Fe<sub>x</sub>Ga<sub>1-x</sub>)Cl<sub>4</sub>, with mixed magnetic and non-magnetic anions  
 SATO A., OJIMA E., AKUTSU H., KOBAYASHI H., KOBAYASHI A., CASSOUX P.  
*Chem. Lett.*, 1998, pp. 673-674 (LCC-5060).

Incorporation of a stilbazole derivative in the hydrogen-bonded chain of L-tartrate : toward a one-step optimization of molecular and bulk second-order nonlinearities

LACROIX P.G., DARAN J.-C., NAKATANI K.

*Chem. Mater.*, 1998, **10**, pp. 1109-1114 (LCC-4826).

Synthesis, crystal structure, and second-order nonlinear optical properties of a new phase-matchable cyanine dye

LACROIX P.G., MALFANT I., PAYRASTRE C., WOLF J.-G., BONVOISIN J., NAKATANI K.

*Chem. Mater.*, 1998, **10**, pp. 1135-1140 (LCC-4827).

New hybrid materials combining nickel bis(dithiolene) metal complexes and reducible stilbazolium cations. A search for possible interplay between optical nonlinearity and conductivity

MALFANT I., CORDENTE N., LACROIX P.G., LEPETIT C.

*Chem. Mater.*, 1998, **10**, pp. 4079-4087 (LCC-5247).

The second example of spin conversion governed by molecular vibrations: a novel ferrous complex resulting from hexacoordination of a Schiff base with a  $N_4O_2$  donor set

BOUSSEKSOU A., SALMON L., VARRET F., TUCHAGUES J.-P.

*Chem. Phys. Lett.*, 1998, **282**, pp. 209-214 (LCC-5113).

Quantum mechanical exchange coupling in polyhydride and dihydrogen complexes

SABO-ETIENNE S., CHAUDRET B.

*Chem. Rev.*, 1998, **98**, pp. 2077-2091 (LCC-5189).

Is alkylation the main mechanism of action of the antimalarial drug artemisinin?

ROBERT A., MEUNIER B.

*Chem. Soc. Rev.*, 1998, **27**, pp. 273-279 (LCC-5061).

Zirconium species as tools in phosphorus chemistry II [ $Cp_2ZrMe_2$ ], a useful reagent

MAJORAL J.-P., IGAU A.

*Coord. Chem. Rev.*, 1998, **176**, pp. 1-16 (LCC-5511).

Zirconocene [ $Cp_2Zr$ ] synthon and benzynezirconocene complexes as tools in main group element chemistry

MAJORAL J.-P., MEUNIER P., IGAU A., PIRIO N., ZABLOCKA M., SKOWRONSKA A., BREDEAU S.

*Coord. Chem. Rev.*, 1998, **178-180**, pp. 145-167 (LCC-5524).

Coordination chemistry of mononuclear iron carbonyl complexes

BRUNET J.-J., CHAUVIN R., DIALLO O.S., KINDELA F.B., LEGLAYE P., NEIBECKER D.

*Coord. Chem. Rev.*, 1998, **178-180**, pp. 331-351 (LCC-5295).

Chemistry of bis(dihydrogen) ruthenium complexes and their derivatives

SABO-ETIENNE S., CHAUDRET B.

*Coord. Chem. Rev.*, 1998, **178-180**, pp. 381-407 (LCC-5281).

Phosphine-terminated dendrimers. Synthesis and complexation properties

CAMINADE A.-M., LAURENT R., CHAUDRET B., MAJORAL J.-P.

*Coord. Chem. Rev.*, 1998, **178-180**, pp. 793-821 (LCC-5207).

The major contribution of phosphorus to the elaboration, the reactivity, and the characterization of giant molecular trees (dendrimers)

CAMINADE A.-M., MAJORAL J.-P.

*ECC*, 1998, **3**, pp. 9-16 (LCC-5208).

The reducing power of  $SmI_2$

CHAUVIN R.

*ECC*, 1998, **3**, pp. 27-28 (LCC-5501).

Activation of dichloromethane by (phosphane)<sub>n</sub> rhodium(I) complexes - X-ray structure of  $\{(\text{PEt}_3)_2\text{RhCl}\}_2(\mu\text{-Cl}_2(\mu\text{-CH}_2))$

BRUNET J.-J., COUILLENS X., DARAN J.-C., DIALLO O.S., LEPETIT C., NEIBECKER D.  
*Eur. J. Inorg. Chem.*, 1998, pp. 349-353 (LCC-4845).

Ethylene polymerization with hydridotris(pyrazol)boratoniobium complexes as precursors

JAFFART J., NAYRAL C., CHOUKROUN R., MATHIEU R., ETIENNE M.

*Eur. J. Inorg. Chem.*, 1998, pp. 425-428 (LCC-4829).

Oxidation of polycyclic aromatic hydrocarbons catalyzed by iron tetrasulfophthalocyanine FePcS : inverse isotope effects and oxygen labeling studies

SOROKIN A., MEUNIER B.

*Eur. J. Inorg. Chem.*, 1998, pp. 1269-1281 (LCC-5064).

The inferior p-donor ability of phosphanyl versus amino substituents : consequences on the stability and reactivity of phosphanyl- and aminocarbenes

GOUMRI S., LERICHE Y., GORNITZKA H., BACEIREDO A., BERTRAND G.

*Eur. J. Inorg. Chem.*, 1998, pp. 1539-1542 (LCC-5209).

A very high-spin molecule : preparation, characterization and magnetic properties of an Fe<sup>III</sup>-Gd<sup>III</sup> complex with an S=12/2 ground state

COSTES J.-P., DUPUIS A., LAURENT J.-P.

*Eur. J. Inorg. Chem.*, 1998, pp. 1543-1546 (LCC-5111).

Synthesis, structure, and physical properties of tetraalkylammonium bis(5,6-dihydro-1,4-dithiin-2,3-diselenolato)nickelate,  $(\text{R}_4\text{N})[\text{Ni}(\text{ddds})_2]$  (R=Me, Et, and *n*Bu), and neutral complex  $[\text{Ni}(\text{ddds})_2]_2$

FUJIWARA H., OJIMA E., KOBAYASHI H., COURCET T., MALFANT I., CASSOUX P.

*Eur. J. Inorg. Chem.*, 1998, pp. 1631-1639 (LCC-5210).

Synthesis of bis[di(2-pyridyl)methyl]amine (BDPMA) by a novel one-pot multi-step reductive amination with molecular sieves and Zn/*i*PrOH

RENZ M., HEMMERT C., MEUNIER B.

*Eur. J. Org. Chem.*, 1998, pp. 1271-1273 (LCC-5065).

Enaminones in the synthesis of new polyaza heterocycles

BEJAN E., AIT-HADDOU H., DARAN J.-C., BALAVOINE G.G.A.

*Eur. J. Org. Chem.*, 1998, pp. 2907-2912 (LCC-5455).

Interplay between chains of S=5/2 localised spins and two-dimensional sheets of organic donors in the synthetically built magnetic multilayer 1-(BETS)<sub>2</sub>FeCl<sub>4</sub>

BROSSARD L., CLERAC R., COULON C., TOKUMOTO M., ZIMAN T., PETROV D.K., LAUKIN V.N., NAUGHTON M., AUDOUARD A., GOZE F., KOBAYASHI A., KOBAYASHI H., CASSOUX P.

*Eur. Phys. J., B*, 1998, **1**, pp. 439-452 (LCC-4830).

On bistability in molecular systems : photomagnetic studies of inorganic complexes

VARRET F., CONSTANT-MACHADO H., DORMANN J.L., GOUJON A., JEFTIC J., NOGUES M., BOUSSEKSOU A., KLOKISHNER S., DOLBECQ A., VERDAGUER M.

*Hyperfine Inter.*, 1998, **113**, pp. 37-46 (LCC-5503).

Homo- (4f, 4f) and heterodimetallic (4f, 4f') complexes. The first structurally characterized example of a heterodimetallic (Yb, La) complex (1'). Magnetic properties of 1' and of a homodinuclear (Gd, Gd) analogue

COSTES J.-P., DAHAN F., DUPUIS A., LAGRAVE S., LAURENT J.-P.

*Inorg. Chem.*, 1998, **37**, pp. 153-155 (LCC-4831).

Influence of hydrogen bonding on the spectroscopic properties and on the reactivity of ruthenium hydrido dihydrogen complexes

GUARI Y., AYLLON J.A., SABO-ETIENNE S., CHAUDRET B., HESSEN B.

*Inorg. Chem.*, 1998, **37**, pp. 640-644 (LCC-4898).

Syntheses, crystal structures, and NLO properties of new chiral inorganic chromophores for second-harmonic generation

LENOBLE G., LACROIX P.G., DARAN J.-C., DI BELLA S., NAKATANI K.

*Inorg. Chem.*, 1998, **37**, pp. 2158-2165 (LCC-4865).

Synthesis, structure, and physical properties of nickel bis(dithiolene) metal complexes, [Ni(dmit)<sub>2</sub>], with highly polar cyanine dyes

MALFANT I., ANDREU R., LACROIX P.G., FAULMANN C., CASSOUX P.

*Inorg. Chem.*, 1998, **37**, pp. 3361-3370 (LCC-5066).

Fluxionality and isomerism of the bis(dihydrogen) complex RuH<sub>2</sub>(H)<sub>2</sub>(PCy)<sub>2</sub>: INS, NMR, and theoretical studies

RODRIGUEZ V., SABO-ETIENNE S., CHAUDRET B., THOBURN J., ULRICH S., LIMBACH H.-H., ECKERT J., BARTHELAT J.-C., HUSSEIN K., MARSDEN C.J.

*Inorg. Chem.*, 1998, **37**, pp. 3475-3485 (LCC-5424).

Preparation of the new bis(phenanthroline) ligand "Clip-Phen" and evaluation of the nuclease activity of the corresponding copper complex

PITIE M., DONNADIEU B., MEUNIER B.

*Inorg. Chem.*, 1998, **37**, pp. 3486-3489 (LCC-5067).

Transition metal derivatives of a ((dimethylamino)ethyl)cyclopentadienyl ligand. Synthesis and structures of amino-containing cyclopentadienyl derivatives of cobalt(I) and -(III) including water-soluble compounds

PHILIPPOPOULOS A.I., BAU R., POILBLANC R., HADJILIADIS N.

*Inorg. Chem.*, 1998, **37**, pp. 4822-4827 (LCC-5211).

Structural and magnetic properties of M(mnt)<sub>2</sub> salts (M=Ni, Pt, Cu) with a ferrocene-based cation, [FcCH<sub>2</sub>N(CH<sub>3</sub>)<sub>3</sub>]<sup>+</sup>. Interplay between M...M and M...S intermolecular interactions

PULLEN A.E., FAULMANN C., POKHODNYA K.I., CASSOUX P., TOKUMOTO M.

*Inorg. Chem.*, 1998, **37**, pp. 6714-6720 (LCC-5212).

New annelated 2,2' : 2"-terpyridines and their Ru(II) complexes : synthesis and characterisation

SASAKI I., DARAN J.-C., AIT-HADDOU H., BALAVOINE G.G.A.

*Inorg. Chem. Commun.*, 1998, **1**, pp. 354-357 (LCC-5305).

Homodinuclear lanthanide complexes : Ln<sub>2</sub>L<sub>3</sub> (H<sub>2</sub>L=tetradentate Schiff bases). Magnetic properties (solid state) and spectroscopic studies (solution)

COSTES J.-P., DUPUIS A., LAURENT J.-P.

*Inorg. Chim. Acta*, 1998, **268**, pp. 125-130 (LCC-4832).

Synthesis and physical studies of copper(I) complexes with the ambidentate ligand (1-piperidyl-2-diphenylphosphino)ethane. X-ray structural characterization of a copper(I) dimer

TURPIN R., BRUNO F., LEGROS J.-P., CASTAN P.

*Inorg. Chim. Acta*, 1998, **271**, pp. 217-221 (LCC-5481).

Sulfonated and acetamidofunctionalized tetraarylporphyrins as biomimetic oxidation catalysts under aqueous conditions

SONG R., ROBERT A., BERNADOU J., MEUNIER B.

*Inorg. Chim. Acta*, 1998, **272**, pp. 228-234 (LCC-4884).

General synthesis of "salicylaldehyde half-unit complexes" : structural determination and use as synthon for the synthesis of dimetallic or trimetallic complexes and of "self-assembling ligand complexes"

COSTES J.-P., DAHAN F., FERNANDEZ FERNANDEZ M.B., FERNANDEZ-GARCIA M.I., GARCIA-DEIBE A., SANMARTIN J.

*Inorg. Chim. Acta*, 1998, **274**, pp. 73-81 (LCC-5059).

- Synthesis, UV-visible and electrochemical studies of lipophilic and hydrophilic lanthanide(III) bis(porphyrinates)<sup>1</sup>  
SPYROULIAS G.A., DE MONTAUZON D., MAISONNAT A., POILBLANC R., COUTSOLELOS A.G.  
*Inorg. Chim. Acta*, 1998, **275-276**, pp. 182-191 (LCC-5431).
- Transition metal complexes with nitronyl nitroxides. Part 1. Crystal structure and properties of square-planar trans radical-copper (II) -radical complexes  
JIANG Z.-H., SUN B.-W., LIAO D.-Z., WANG G.-L., DAHAN F., TUCHAGUES J.-P.  
*Inorg. Chim. Acta*, 1998, **279**, pp. 69-75 (LCC-5083).
- Transition metal complexes with nitronyl nitroxides. Part 2. Crystal structure and magnetism of manganese(II) compounds including radicals and their reduced derivatives  
JIANG Z.-H., SUN B.-W., LIAO D.-Z., WANG G.-L., DONNADIEU B., TUCHAGUES J.-P.  
*Inorg. Chim. Acta*, 1998, **279**, pp. 76-84 (LCC-5084).
- Electrocrystallization and characterization of phosphonium-based radical anion salts [MePh<sub>3</sub>P][Pd(dmit)<sub>2</sub>]<sub>3</sub> and [Ph<sub>4</sub>P][Pd(dmit)<sub>2</sub>]<sub>3</sub> (dmit=1,3-dithiole-2-thione-4,5-dithiolate). Comparison with related Pd and Ni isologues  
PULLEN A.E., FAULMANN C., LIU H.L., TANNER D.B., ABOUD K.A., REYNOLDS J.R.  
*Inorg. Chim. Acta*, 1998, **282**, pp. 90-95 (LCC-5213).
- Organometallic Rh(I) and Rh(III) complexes with [2-(dimethylamino)ethyl] cyclopentadiene  
PHILIPPOPOULOS A.I., POILBLANC R., HADJILIADIS N.  
*Inorg. Chim. Acta*, 1998, **283**, pp. 24-29 (LCC-5509).
- Zirconocene-ate Zwitterionic complexes : preparation and X-ray structure  
MIQUEL Y., IGAU A., DONNADIEU B., MAJORAL J.-P., PIRIO N., MEUNIER P.  
*J. Am. Chem. Soc.*, 1998, **120**, pp. 3504-3505 (LCC-4864).
- Phosphorus-containing dendrimers : chemoselective functionalization of internal layers  
LARRE C., DONNADIEU B., CAMINADE A.-M., MAJORAL J.-P.  
*J. Am. Chem. Soc.*, 1998, **120**, pp. 4029-4030 (LCC-4836).
- Exchange couplings between a hydride and a stretched dihydrogen ligand in ruthenium complexes  
GUARI Y., SABO-ETIENNE S., CHAUDRET B.  
*J. Am. Chem. Soc.*, 1998, **120**, pp. 4228-4229 (LCC-4899).
- NMR and INS line shapes of transition metal hydrides in the presence of coherent and incoherent dihydrogen exchange  
LIMBACH H.-H., ULRICH S., GRUNDEMANN S., BUNTKOWSKI G., SABO-ETIENNE S., CHAUDRET B., KUBAS G.J., ECKERT J.  
*J. Am. Chem. Soc.*, 1998, **120**, pp. 7929-7943 (LCC-5194).
- Efficient oxidation of 2'-deoxyguanosine by Mn-TMPyP/KHSO<sub>5</sub> to imidazolone dIz without formation of 8-oxo-dG  
VIALAS C., PRATVIEL G., CLAPAROLS C., MEUNIER B.  
*J. Am. Chem. Soc.*, 1998, **120**, pp. 11548-11553 (LCC-5214).
- Chemistry within megamolecules : regiospecific functionalization after construction of phosphorus dendrimers  
LARRE C., BRESSOLLES D., TURRIN C.O., DONNADIEU B., CAMINADE A.-M., MAJORAL J.-P.  
*J. Am. Chem. Soc.*, 1998, **120**, pp. 13070-13082 (LCC-5215).
- Two new cation radical bis (ethylenedithio)tetraselenafulvalene (BETS) salts,  $\alpha$ -(BETS)<sub>6</sub>Bi<sub>3</sub>Cl<sub>12</sub>.PhCl and (BETS)<sub>2</sub>Bi<sub>2</sub>Cl<sub>8</sub> : synthesis, structure and conducting properties. First characterisation of the new trinuclear anion [Bi<sub>3</sub>Cl<sub>12</sub>]<sup>3-</sup>  
KUSHCH N.D., DYACHENKO O.A., GRITSENKO V.V., CASSOUX P., FAULMANN C., KOBAYASHI A., KOBAYASHI H.  
*J. Chem. Soc., Dalton Trans.*, 1998, pp. 683-688 (LCC-4833).

An original heterodinuclear VO<sup>2+</sup>, Gd<sup>3+</sup> complex with a nonet ground state

COSTES J.-P., DUPUIS A., LAURENT J.-P.

*J. Chem. Soc., Dalton Trans.*, 1998, pp. 735-736 (LCC-4835).

Synthesis and structural characterization of dialkyldithiocarbamate complexes of niobium

FERNANDEZ-BAEZA J., JALON F.A., OTERO A., RODRIGO-BLANCO M.-E., ETIENNE M.

*J. Chem. Soc., Dalton Trans.*, 1998, pp. 769-773 (LCC-4834).

New intermediate spin chloro-iron(III) complexes with a mixed nitrogen sulfur coordination sphere

KOPF M.-A., VARECH D., TUCHAGUES J.-P., MANSUY D., ARTAUD I.

*J. Chem. Soc., Dalton Trans.*, 1998, pp. 991-998 (LCC-5114).

Bridging ability of a novel polydentate ligand (H<sub>2</sub>L) comprising an oxime function. Structures of a mononuclear precursor [NiL] and a dinuclear Cu<sup>II</sup><sub>2</sub> complex. Magnetic properties of mononuclear (Ni<sup>II</sup> and Cu<sup>II</sup>), dinuclear (Cu<sup>II</sup><sub>2</sub>, Ni<sup>II</sup><sub>2</sub>, Ni<sup>II</sup>Cu<sup>II</sup> and Cu<sup>II</sup>Cr<sup>III</sup>) and trinuclear (Cu<sup>II</sup><sub>3</sub>, Cu<sup>II</sup>Mn<sup>II</sup>Cu<sup>II</sup> and Cu<sup>II</sup>Zn<sup>II</sup>Cu<sup>II</sup>) complexes

COSTES J.-P., DAHAN F., DUPUIS A., LAURENT J.-P.

*J. Chem. Soc., Dalton Trans.*, 1998, pp. 1307-1314 (LCC-5112).

Synthesis, crystal structure and electrochemical properties of [NBu<sub>4</sub>][Ni(mdt)<sub>2</sub>]: a potential precursor for new materials (mdt = 1,3-dithiol-4,5-dithiolate)

VELDHUIZEN Y.S.J., VELDMAN N., SPEK A.L., CASSOUX P., CARLIER R., MULDER M.J.J., HAASNOOT J.G., REEDIJK J.

*J. Chem. Soc., Dalton Trans.*, 1998, pp. 2989-2993 (LCC-5416).

Synthesis and characterization of molybdenum based colloidal particles

MORENO B., VIDONI O., OVALLES C., CHAUDRET B., URBINA C., KRENTZIEN H.

*J. Colloid Interf. Sci.*, 1998, **207**, pp. 251-257 (LCC-5282).

Transition metal complexes and cycloaddition products of pentafluorophenyl isocyanide

LENTZ D., ANIBARRO M., PREUGSCHAT D., BERTRAND G.

*J. Fluor. Chem.*, 1998, **89**, pp. 73-81 (LCC-5180).

Crystal structure at 180°K of bis-3,5-diisopropylsalicylatobisdimethylsulfoxidozinc(II) and the inhibition of seizures and polymorphonuclear leukocyte chemoluminescence

MORGANT G., VIOSSAT B., DARAN J.-C., ROCH-ARVEILLER M., GIROUD J.-P., DUNG N.-H., SORENSON J.R.J.

*J. Inorg. Biochem.*, 1998, **70**, pp. 137-143 (LCC-5458).

New convenient one-pot methods of conversion of alkynes to cyclobutenediones or  $\alpha$ ,  $\beta$ -unsaturated carboxylic acids using novel reactive iron carbonyl reagents

PERIASAMY M., RAMESHKUMAR C., RADHAKRISHNAN U., BRUNET J.-J.

*J. Org. Chem.*, 1998, **63**, pp. 4930-4935 (LCC-5297).

Study of cyclic borinates obtained from piperidine- and piperazine alcohols by spectroscopic methods and X-ray crystallography

HOPFL H., FARFAN N., CASTILLO D., SANTILLAN R., GUTIERREZ A., DARAN J.-C.

*J. Organomet. Chem.*, 1998, **553**, pp. 221-239 (LCC-5457).

Synthesis, characterization and electrochemical studies of the heterometallic diclusters [ $\{\text{Fe}_2(\mu\text{-CO})(\text{CO})_6(\mu\text{-PPh}_2)\text{Au}\}_2(\text{diphosphine})$ ] (diphosphine = dpmp, dppip, dppe, dppp)

FERRER M., JULIA A., REINA R., ROSSELL O., SECO M., DE MONTAUZON D.

*J. Organomet. Chem.*, 1998, **560**, pp. 147-153 (LCC-5434).

Synthesis and X-ray crystal structure of bis(germyl)carbodiimides. Spectroscopic characterization of (hydrogeno) (germyl) cyanimides

DAHROUCH M., RIVIERE-BAUDET M., GORNITZKA H., BERTRAND G.

*J. Organomet. Chem.*, 1998, **562**, pp. 191-195 (LCC-5159).



- Mono and difunctionalisation of chiral ferrocenyl bis-acetals. X-ray crystal structure of bis-1,1'-[2S, 4S)-(hydroxymethyl)-2-dioxane 1,3]-ferrocene  
 IFTIME G., DARAN J.-C., MANOURY E., BALAVOINE G.G.A.  
*J. Organomet. Chem.*, 1998, **565**, pp. 115-124 (LCC-5459).
- New chiral zwitterionic iron hydride complexes with ephosium" and "valphosium" ligands  
 BRUNET J.-J., CHAUVIN R., CHIFFRE J., HUGUET S., LEGLAYE P.  
*J. Organomet. Chem.*, 1998, **566**, pp. 117-123 (LCC-5298).
- Reaction of aminocarbene complexes of chromium with alkynes 10. From large to small cyclic amines : single versus double alkyne insertions  
 LAFOLLEE-BEZZENINE S., PARLIER A., RUDLER H., VAISSERMANN J., DARAN J.-C.  
*J. Organomet. Chem.*, 1998, **567**, pp. 83-100 (LCC-5460).
- Selective synthesis of ferrocenes  
 BALAVOINE G.G.A., DARAN J.-C., IFTIME G., MANOURY E., MOREAU-BOSSUET C.  
*J. Organomet. Chem.*, 1998, **567**, pp. 191-198 (LCC-5456).
- Synthesis and reactivity of  $[(R^1R^2N)_2PH]Fe(CO)_4$  complexes. X-ray crystal structure of  $[(Ph_2N)_2PH]Fe(CO)_4$   
 BRUNET J.-J., CHAUVIN R., DIALLO O.S., DONNADIEU B., JAFFART J., NEIBECKER D.  
*J. Organomet. Chem.*, 1998, **570**, pp. 195-200 (LCC-5343).
- Improved synthesis of  $KHCr(CO)_5$  and comparative coordination chemistry from  $KHCr(CO)_5$  and  $KHFe(CO)_4$   
 BRUNET J.-J., CHAUVIN R., DONNADIEU B., LEGLAYE P., NEIBECKER D.  
*J. Organomet. Chem.*, 1998, **571**, pp. 7-13 (LCC-5299).
- Surface chemistry of colloidal platinum : vibrational coupling in the infrared spectrum of CO adsorbed on colloidal platinum in liquid dispersion  
 DE CARO D., BRADLEY J.S.  
*Langmuir*, 1998, **14**, pp. 245-247 (LCC-5485).
- Wide angles X-ray scattering (W.A.X.S.) and H.R.E.M. Studies on nanoscale cobalt and cobalt oxide colloids  
 VERELST M., SNOECK E., OULD ELY T., LECANTE P., AMIENS C., ROUCAU C., MOSSET A., CHAUDRET B.  
*Mater. Sci. Forum*, 1998, **269-272**, pp. 403-408 (LCC-5068).
- Elaboration, characterization and magnetic properties of cobalt fine particles  
 RESPAUD M., BROTO J.-M., THOMAS L., BARBARA B., RAKOTO H., GOIRAN M., FERT A., SNOECK E., VERELST M., LECANTE P., OSUNA J., OULD ELY T., AMIENS C., CHAUDRET B.  
*Mater. Sci. Forum*, 1998, **269-272**, pp. 949-954 (LCC-4856).
- Optical detection of the spin transition by reflectivity: application to  $[Fe_xCo_{1-x}](btr)_2(NCS)_2 \cdot H_2O$   
 MORSCHEIDT W., JEFTIC J., CODJOVI E., LINARES J., BOUSSEKSOU A., CONSTANT-MACHADO H., VARRET F.  
*Meas. Sci. Technol.*, 1998, **9**, pp. 1311-1315 (LCC-5115).
- Bis(ethylenedithio)tetraselenafulvalene : short-cut synthesis, x-ray crystal structure and p-electron density distribution  
 COURCET T., MALFANT I., POKHODNIA K., CASSOUX P.  
*New J. Chem.*, 1998, pp. 585-589 (LCC-5161).
- Transition-metal derivatives of the functionalized cyclopentadienyl ligand. XVI. Synthesis of the bridged complexes  $[(\mu-h^5-C_5I_4PPh_2)M(CO)_2]_2$  (M=Cr, Mo, W). X-ray crystal structure of the dihydride derivative  $[(\mu-h^5-C_5H_4PPh_2)W(CO)_2H]_2$   
 BRUMAS-SOULA B., DAHAN F., POILBLANC R.  
*New J. Chem.*, 1998, **22**, pp. 15-23 (LCC-4837).

Sequential addition of H<sub>2</sub>O<sub>2</sub>, pH and solvent effects as key factors in the oxidation of 2,4,6-trichlorophenol catalyzed by iron tetrasulfophthalocyanine  
HADASCH A., SOROKIN A., RABION A., MEUNIER B.  
*New J. Chem.*, 1998, **22**, pp. 45-51 (LCC-4849).

Synthesis and characterization of two ruthenium(II) heteroleptic complexes of annelated derivatives of 2,2'-bipyridine  
AMOUYAL E., PENAUD-BERRUYER F., AZHARI D., AIT-HADDOU H., FONTENAS C., BEJAN E., DARAN J.-C., BALAVOINE G.G.A.  
*New J. Chem.*, 1998, **22**, pp. 373-380 (LCC-4858).

Platinum nanoparticles stabilized by CO and octanethiol ligands or polymers : FT-IR, NMR, HREM and WAXS studies  
DASSENROY F., PHILIPPOT K., OULD ELY T., AMIENS C., LECANTE P., SNOECK E., MOSSET A., CASANOVE M.-J., CHAUDRET B.  
*New J. Chem.*, 1998, **22**, pp. 703-711 (LCC-5190).

Transition-metal derivatives of the functionalized cyclopentadienyl ligand. XVIII. Syntheses and structures of a series of tetranuclear cyclic complexes  $\{[(CO)_3M(\mu, \eta^5-C_5H_4PPh_2)]_2M'M''\}$  (M=Cr, Mo or W, M'=M''=Ag; M=Cr or Mo, M'=M''=Au; M=Mo, M'=Ag, M''=Au). Isotopomerism and NMR evidence of a double epimerization process  
BRUMAS-SOULA B., DAHAN F., POILBLANC R.  
*New J. Chem.*, 1998, **22**, pp. 1067-1074 (LCC-5216).

Synthesis, crystal structure and INDO calculation of the molecular hyperpolarizability of a highly polarizable merocyanine dye  
LACROIX P.G., DARAN J.-C., CASSOUX P.  
*New J. Chem.*, 1998, **22**, pp. 1085-1091 (LCC-5217).

Investigation of the surface structure of colloidal platinum by infrared spectroscopy of adsorbed CO  
DE CARO D., BRADLEY J.S.  
*New J. Chem.*, 1998, **22**, pp. 1267-1273 (LCC-5484).

Use of vanadocene as an organometallic reducing agent for the preparation of ultrafine magnetic iron powder and organized iron particles from iron salts  
CHOUKROUN R., DE CARO D., MATEO S., AMIENS C., CHAUDRET B., SNOECK E., RESPAUD M.  
*New J. Chem.*, 1998, **22**, pp. 1295-1299 (LCC-5218).

Structural studies of a dinuclear (Cu, Gd) and two trinuclear (Cu<sub>2</sub>, Ln) complexes (Ln=Ce, Er). Magnetic properties of two original (Cu, Gd) complexes  
COSTES J.-P., DAHAN F., DUPUIS A., LAURENT J.-P.  
*New J. Chem.*, 1998, **22**, pp. 1525-1529 (LCC-5219).

Synthesis and biodistribution of new oxo and nitrido <sup>99m</sup>Tc complexes with asymmetrical potentially dianionic or trianionic tetradentate SNNO ligands derived from methyl-2-aminocyclopentene-1-dithiocarboxylic acid  
OUHAYOUN E., CROS G., DARBIEU M.-H., TAFANI J.A.M., FABRE J., ESQUERRE J.-P., COULAIS Y., BELHADJ-TAHAR H.  
*Nucl. Med. Biol.*, 1998, **25**, pp. 65-69 (LCC-5116).

Neutral and cationic tetracoordinated aluminum complexes featuring tridentate nitrogen donors : synthesis, structure, and catalytic activity for the ring-opening polymerization of propylene oxide and (D, L)-lactide  
EMIG N., NGUYEN H., KRAUTSCHEID H., REAU R., CAZAUX J.-B., BERTRAND G.  
*Organometallics*, 1998, **17**, pp. 3599-3608 (LCC-5160).

Hydride and dihydrogen ruthenium complexes of the tripyrrolylphosphine ligand  
RODRIGUEZ V., DONNADIEU B., SABO-ETIENNE S., CHAUDRET B.  
*Organometallics*, 1998, **17**, pp. 3809-3814 (LCC-5195).

Electrophilic additions of mercury and gold species to the anion carbide cluster  $[\text{Fe}_3\text{C}(\text{CO})_{14}]^{2-}$ . X-ray crystal structures of  $(\text{NEt}_4)_5\text{Fe}_5\text{C}(\text{CO})_{14}\{\mu\text{-HgW}(\text{CO})_3\text{Cp}\}$  and  $(\text{NEt}_4)_2[\mu_4\text{-Hg}\{\text{Fe}_5\text{C}(\text{CO})_{14}\}_2]$   
REINA R., RIBA O., ROSSELL O., SECO M., GOMEZ-SAL P., MARTIN A., DE MONTAUZON D., MARI A.

*Organometallics*, 1998, **17**, pp. 4127-4135 (LCC-5435).

Stoichiometric and catalytic activation of allyldimethylsilane. Synthesis of  $[\text{RuH}_2\{\text{h}^4\text{-HSiMe}_2(\text{CH}=\text{CHMe})\}(\text{PCy}_3)_2]$

DELPECH F., SABO-ETIENNE S., DONNADIEU B., CHAUDRET B.

*Organometallics*, 1998, **17**, pp. 4926-4928 (LCC-5283).

New bis-biphosphole - transition-metal complexes : spontaneous isomerization of trans-meso- into cis-0-  
[Ru(trifluoroacetate)<sub>2</sub> (1,1'-diphenyl-3,3',4,4'- tetramethyl-2,2'-biphosphole)<sub>2</sub>]

TISSOT O., GOUYGOU M., DARAN J.-C., BALAVOINE G.G.A.

*Organometallics*, 1998, **17**, pp. 5927-5930 (LCC-5306).

Synthesis of 1,n-bis (2,5-diphenylphosphol-1-yl)alkanes

BOUSQUET S., BRUNET J.-J., COURCET T., NEIBECKER D.

*Phosphorus Sulfur Silicon Relat. Elem.*, 1998, **142**, pp. 117-124 (LCC-5300).

Surface effects on the magnetic properties of ultrafine cobalt particles

RESPAUD M., BROTO J.-M., RAKOTO H., FERT A., THOMAS L., BARBARA B., VERELST M., SNOECK E., LECANTE P., MOSSET A., OSUNA J., OULD ELY T., AMIENS C., CHAUDRET B.

*Phys. Rev. B-Cond. Matt.*, 1998, **57**, pp. 2925-2935 (LCC-4897).

Coupling of metal-insulator and antiferromagnetic transitions in the highly correlated organic conductor incorporating magnetic anions, 1-BETS<sub>2</sub>FeBr<sub>x</sub>Cl<sub>4-x</sub> [BETS = bis(ethylenedithio)tetraselenafulvalene]

AKUTSU H., KATO K., OJIMA E., KOBAYASHI H., TANAKA H., KOBAYASHI A., CASSOUX P.

*Phys. Rev. B-Cond. Matt.*, 1998, **58**, pp. 9294-9302 (LCC-5417).

Magnetization measurements on fine cobalt particles

RESPAUD M., BROTO J.-M., RAKOTO H., OUSSET J.C., OSUNA J., OULD ELY T., AMIENS C., CHAUDRET B., ASKENAZY S.

*Physica B*, 1998, **246-247**, pp. 532-536 (LCC-5191).

High field ferromagnetic resonance in fine particles

RESPAUD M., GOIRAN M., YANG H., BROTO J.-M., OULD ELY T., AMIENS C., CHAUDRET B., ASKENAZY S.

*Physica B*, 1998, **246-247**, pp. 580-583 (LCC-5192).

Palladium complexes of unsaturated secondary aziridines. Crystal structure of trans-dichlorobis (2,2-dimethyl-3-allylaziridine) palladium(II)

BEN CHEIKH R., CHAABOUNI R., DAHAN F., BONNET M.C.

*Polyhedron*, 1998, **17**, pp. 185-192 (LCC-4818).

Isolation of a benzenze valence isomer with one-electron phosphorus-phosphorus bonds

CANAC Y., BOURISSOU D., BACEIREDO A., GORNITZKA H., SCHOELLER W., BERTRAND G.

*Science*, 1998, **279**, pp. 2080-2082 (LCC-4853).

A coupled metal-insulator and antiferromagnetic transition of 1-(BETS)<sub>2</sub>FeCl<sub>4</sub> under high-pressure and magnetic field [BETS = bis(ethylenedithio)tetraselenafulvalene]

AKUTSU H., KATO K., ARAI E., KOBAYASHI H., KOBAYASHI A., TOKUMOTO M., BROSSARD L., CASSOUX P.

*Solid State Commun.*, 1998, **105**, pp. 485-489 (LCC-4838).

The reaction of enamines with Grignard reagents : synthesis of  $\alpha$ ,  $\beta$ -unsaturated ketones

FONTENAS C., AIT-HADDOU H., BEJAN E., BALAVOINE G.G.A.

*Synth. Commun.*, 1998, **28**, pp. 1743-1753 (LCC-4859).

Push-pull azulene-based chromophores with nonlinear optical properties

IFTIME G., LACROIX P.G., NAKATANI K., RAZUS A.C.

*Tetrahedron Lett.*, 1998, **39**, pp. 6853-6856 (LCC-5250).

"Methyldiopium" and "methylbinapium", chiral phosphonium-phosphine ligands

LEGLAYE P., DONNADIEU B., BRUNET J.-J., CHAUVIN R.

*Tetrahedron Lett.*, 1998, **39**, pp. 9179-9182 (LCC-5301).

Divergent approaches to phosphorus-containing dendrimers and their functionalization

MAJORAL J.-P., CAMINADE A.-M.

*Top. Curr. Chem.*, 1998, **197**, pp. 79-124 (LCC-5069).