

Publications LCC 2000

A new precursor for the synthesis of porphyrazine : 2,3-bis(4-methylphenyl)maleonitrile

FAULMANN C., CASSOUX P., PULLEN A.E.

Acta Crystallogr., Sect. C, 2000, **C56**, pp. 384-385 (LCC-5352).

MALDI TOF mass spectrometry for the characterization of phosphorus-containing dendrimers. Scope and limitations

BLAIS J.-C., TURRIN C.O., CAMINADE A.-M., MAJORAL J.-P.

Anal. Chem., 2000, **72**, pp. 5097-5105 (LCC-5353).

The assembly of well-defined nano building blocks yields novel mesostructured hybrid compounds

SOLLER-ILIA G.J., ROZES L., BOGGIANO M.K., SANCHEZ C., TURRIN C.O., CAMINADE A.-M., MAJORAL J.-P.

Angew. Chem. (Int. Ed. Engl.), 2000, **39**, pp. 4249-4254 (LCC-5355).

Early transition metal α -diazoalkane complexes

CADIERNO V., ZABLOCKA M., DONNADIEU B., IGAU A., MAJORAL J.-P.

Angew. Chem. (Int. Ed. Engl.), 2000, **39**, pp. 4524-4528 (LCC-5523).

In vitro and in vivo potentiation of artemisinin and synthetic endoperoxide antimalarial drugs by metalloporphyrins

BENOIT-VICAL F., ROBERT A., MEUNIER B.

Antimicrob. Agents Chemother., 2000, **44**, pp. 2836-2841 (LCC-5356).

Synthesis and use of a novel SnO₂ nanomaterial for gas sensing

NAYRAL C., VIALA E., COLLIERE V., FAU P., SENOCQ F., MAISONNAT A., CHAUDRET B.

Appl. Surf. Sci., 2000, **164**, pp. 219-226 (LCC-5357).

Conceptualisation of diagnostic agents : from empirical in vivo screening to rational in vitro predictive parameters

BOUMAHDI R., DARBIEU M.-H., BELHADJ-TAHAR H.

ATLA, 2000, **28**, pp. 303-314 (LCC-5448).

Oxidative damage generated by an oxo-metalloporphyrin onto human telomeric sequence

VIALAS C., PRATVIEL G., MEUNIER B.

Biochem., 2000, **39**, pp. 9514-9522 (LCC-5358).

Targeting the DNA cleavage activity of copper phenanthroline and clip-phen to A-T tracts via linkage to a poly-N-methylpyrrole

PITIE M., VAN HORN J.D., BRION D., BURROWS C.J., MEUNIER B.

Bioconjugate Chem., 2000, **11**, pp. 892-900 (LCC-5359).

Single versus double nucleophilic addition carboxylic acid enolates to anisole tricarbonyl chromium complexes with formation of arylcarboxylic acids and/or tetrahydrobenzofuran-2,5-diones : the size of the substituents on the acid makes the difference

RUDLER H., COMTE V., BELLASSOUED M., GARRIER E., VAISSERMANN J., DARAN J.-C.

C. R. Acad. Sci., Paris, Sér.II c, 2000, **3**, pp. 113-122 (LCC-5469).

The influence of crystal packing on second-order non-linear optical properties of new chiral ferrocenyl materials

IFTIME G., BALAVOINE G.G.A., DARAN J.-C., LACROIX P.G., MANOURY E.

C. R. Acad. Sci., Paris, Sér.II c, 2000, **3**, pp. 139-146 (LCC-5438).

Oxidation of thiureas with potassium monopersulfate : an efficient method for the environmentally safe degradation

FELL R.T., MEUNIER B.

C. R. Acad. Sci., Paris, Sér.II c, 2000, **3**, pp. 285-288 (LCC-5360).

Application of conversion coatings to the growth of TTF-TCNQ thin films by CVD and conducting nanowires by dipping process

DE CARO D., SAKAH J., BASSO-BERT M., FAULMANN C., LEGROS J.-P., ONDARCUHU T., JOACHIM C., ARIES L., VALADE L., CASSOUX P.

C. R. Acad. Sci., Paris, Sér.II c, 2000, **3**, pp. 675-680 (LCC-5361).

Second generation of a polypyridine ligand to mimic enzymes containing non-heme iron centers
HEMMERT C., MAESTRIN A.P., RENZ M., GORNITZKA H., MEUNIER B.

C. R. Acad. Sci., Paris, Sér.II c, 2000, **3**, pp. 735-741 (LCC-5362).

Origin of the oxygen atom in metalloporphyrin-catalyzed epoxidations with LiOCl as oxidant
ROBERT A., MEUNIER B.

C. R. Acad. Sci., Paris, Sér.II c, 2000, **3**, pp. 771-775 (LCC-5385).

Electrogenerated poly(dendrimers) containing conjugated poly(thiophene) chains

SEBASTIAN R.M., CAMINADE A.-M., MAJORAL J.-P., LEVILLAIN E., HUCHET L., RONCALI J.

Chem. Commun., 2000, pp. 507-508 (LCC-5364).

X-ray crystal structure of the bis(dihydrogen) complex $\text{RuH}_2(\text{H}_2)_2(\text{PCy}_3)_2$

BOROWSKI A.F., DONNADIEU B., DARAN J.-C., SABO-ETIENNE S., CHAUDRET B.

Chem. Commun., 2000, pp. 543-544 (LCC-5425).

Two photochemical pathways in competition : matrix isolation, time-resolved and NMR studies of *cis*- $[\text{Ru}(\text{PMe}_3)_4(\text{H})_2]$

MONTIEL-PALMA V., PERUTZ R.N., GEORGE M.W., JINA O.S., SABO-ETIENNE S.

Chem. Commun., 2000, pp. 1175-1176 (LCC-5426).

$\text{Li}[\text{Cp}_2\text{Zr}(\text{C}=\text{CPh})(\text{h}^2\text{-}1,2\text{-PhC}_2\text{C}=\text{CPh})]$: an anionic zirconium(II) intermediate for carbon-carbon coupling

CHOUKROUN R., ZHAO J.S., LORBER C., CASSOUX P., DONNADIEU B.

Chem. Commun., 2000, pp. 1511-1512 (LCC-5365).

DFT exploration of structural and magnetic properties of [N]annulene ring carbomers

GODARD C., LEPETIT C., CHAUVIN R.

Chem. Commun., 2000, pp. 1833-1834 (LCC-5491).

Gold nanoparticles from self-assembled gold(I) amine precursors

GOMEZ S., PHILIPPOT K., COLLIERE V., CHAUDRET B., SENOCQ F., LECANTE P.

Chem. Commun., 2000, pp. 1945-1946 (LCC-5422).

Unexpected formal [1+3] cycloadditions between azides and σ -zirconated phosphanes : a route to unprecedented phosphazide and iminophosphorane complexes

CADIerno V., ZABLOCKA M., DONNADIEU B., IGAU A., MAJORAL J.-P., SKOWRONSKA A.

Chem. Eur. J., 2000, **6**, pp. 345-352 (LCC-5520).

Naked Au_{55} clusters : dramatic effect of a thiol-terminated dendrimer

SCHMID G., MEYER-ZAIKA W., PUGIN R., SAWITOWSKI T., MAJORAL J.-P., CAMINADE A.-M., TURRIN C.O.

Chem. Eur. J., 2000, **6**, pp. 1693-1697 (LCC-5366).

Azo-azulene derivatives as second-order nonlinear optical chromophores

LACROIX P.G., MALFANT I., IFTIME G., RAZUS A.C., NAKATANI K., DELAIRE J.

Chem. Eur. J., 2000, **6**, pp. 2599-2608 (LCC-5439).

Synthesis of tin and tin oxide nanoparticles of low size dispersity for application in gas sensing

NAYRAL C., VIALA E., FAU P., SENOCQ F., JUMAS J.-C., MAISONNAT A., CHAUDRET B.

Chem. Eur. J., 2000, **6**, pp. 4082-4090 (LCC-5367).

[Cp₂V] migration along an octatetrayne chain : from the monometallic complex [Cp₂V(3-4*h*-*t*BuC=C-C₂-C=CC=C*t*Bu)] to the dimetallic complex [(Cp₂V)₂(1-2*n*:7-8-*t*BuC₂-C=CC=CC₂*t*Bu)]
CHOUKROUN R., DONNADIEU B., LORBER C., PELLNY P.M., BAUMANN W., ROSENTHAL U.
Chem. Eur. J., 2000, **6**, pp. 4505-4509 (LCC-5368).

CVD-grown thin films of molecule-based magnets
DE CARO D., BASSO-BERT M., SAKAH J., CASELLAS H., LEGROS J.-P., VALADE L., CASSOUX P.
Chem. Mater., 2000, **12**, pp. 587-589 (LCC-5369).

Theoretical investigation of the effect of a spin transition on the second-order molecular hyperpolarizability of a bis(salicylaldiminato)Fe^{II} Schiff base complex
AVERSENG F., LEPETIT C., LACROIX P.G., TUCHAGUES J.-P.
Chem. Mater., 2000, **12**, pp. 2225-2229 (LCC-5440).

Organic-inorganic hybrid materials incorporating phosphorus-containing dendrimers
TURRIN C.O., MARAVAL V., CAMINADE A.-M., MAJORAL J.-P., MEHDI A.
Chem. Mater., 2000, **12**, pp. 3848-3856 (LCC-5370).

Raman spectroscopy of the high- and low-spin states of the spin crossover complex Fe(phen)₂(NCS)₂ : an initial approach to estimation of vibrational contributions to the associated entropy change
BOUSSEKSOU A., MCGARVEY J.J., VARRET F., REAL J.A., TUCHAGUES J.-P., DENNIS A.C., BOILLOT M.L.
Chem. Phys. Lett., 2000, **318**, pp. 409-416 (LCC-5504).

BETS as a source of molecular magnetic superconductors (BETS = bis(ethylenedithio)tetraselenafulvalene)
KOBAYASHI H., KOBAYASHI A., CASSOUX P.
Chem. Soc. Rev., 2000, **29**, pp. 325-333 (LCC-5371).

Preparation and antimalarial activities of "trioxaquines", new modular molecules with a trioxane skeleton linked to a 4-aminoquinoline
DECHY-CABARET O., BENOIT-VICAL F., ROBERT A., MEUNIER B.
ChemBioChem, 2000, pp. 281-283 (LCC-5363).

Photocatalytic degradation of atrazine by porphyrin and phthalocyanine complexes
HEQUET V., LE CLOIREC P., GONZALEZ C., MEUNIER B.
Chemosphere, 2000, **41**, pp. 379-386 (LCC-5499).

CVD of Molecule-based Conductors and Magnets
VALADE L., DE CARO D., CASELLAS H., BASSO-BERT M., FAULMANN C., LEGROS J.-P., CASSOUX P.
Electrochemical Society Proceedings, 2000, **13**, pp. 781-786 (LCC-6665).

2-*H*-1,2-azaphosphindoles - synthesis and characterization
CADIERNO V., DONNADIEU B., IGAU A., MAJORAL J.-P.
Eur. J. Inorg. Chem., 2000, pp. 417-421 (LCC-5521).

Zwitterionic organometallates
CHAUVIN R.
Eur. J. Inorg. Chem., 2000, pp. 577-591 (LCC-5493).

Selective and environmentally benign aerobic catalytic oxidation of alcohols by a molybdenum-copper system
LORBER C., SMIDT S.P., OSBORN J.A.
Eur. J. Inorg. Chem., 2000, pp. 655-658 (LCC-5487).

Alkyne[hydrotris(pyrazolyl)borato]tantalum complexes - an ethyl group is a better α -agostic donor than a methyl group
HIERSO J.-C., ETIENNE M.
Eur. J. Inorg. Chem., 2000, pp. 839-842 (LCC-5506).

Pentacarbonylhydrido chromates $M^+[HCr(CO)_5]^-$: reactivity in organic synthesis and homogeneous catalysis

BRUNET J.-J.

Eur. J. Inorg. Chem., 2000, pp. 1377-1390 (LCC-5327).

Metal-mediated oxidation of tertiary alcohols and related fragmentations

WIETZERBIN K., BERNADOU J., MEUNIER B.

Eur. J. Inorg. Chem., 2000, pp. 1391-1406 (LCC-5372).

The ligand 1,10-phenanthroline-2,9-dicarbaldehyde dioxime can act both as a tridentate and as a tetradentate ligand - Synthesis, characterization and crystal structures of its transition metal complexes

ANGELOFF A., DARAN J.-C., BERNADOU J., MEUNIER B.

Eur. J. Inorg. Chem., 2000, pp. 1985-1996 (LCC-5373).

New tetrathiafulvalene-p-spacer-acceptor derivatives : synthesis, crystal structure, optical and electrochemical properties

ANDREU R., MALFANT I., LACROIX P.G., CASSOUX P.

Eur. J. Org. Chem., 2000, pp. 737-741 (LCC-5374).

Synthesis of new soluble annelated polypyridines

AIT-HADDOU H., FONTENAS C., BEJAN E., DARAN J.-C., BALAVOINE G.G.A.

Eur. J. Org. Chem., 2000, pp. 987-994 (LCC-5470).

A new class of ferrocene-based 1,2-bis(phosphanes) possessing only planar chirality

ARGOUARCH G., SAMUEL O., RIAN T O., DARAN J.-C., KAGAN H.

Eur. J. Org. Chem., 2000, pp. 2893-2899 (LCC-5471).

Michael-type addition of amines to the vinyl core of dendrons - Application to the synthesis of multidendritic systems

MARAVAL V., LAURENT R., MERINO S., CAMINADE A.-M., MAJORAL J.-P.

Eur. J. Org. Chem., 2000, pp. 3555-3568 (LCC-5375).

Dynamic triggering of a spin-transition by a pulsed magnetic field

BOUSSEKSOU A., NEGRE N., GOIRAN M., SALMON L., TUCHAGUES J.-P., BOILLOT M.L., BOUKHEDDADEN K.

Eur. Phys. J., B, 2000, **13**, pp. 451-456 (LCC-5376).

Influence of anionic ligands (X) on the nature and magnetic properties of dinuclear $LCuGdX_3 \cdot nH_2O$ complexes (LH₂ standing for tetradentate Schiff base ligands deriving from 2-hydroxy-3-methoxybenzaldehyde and X being Cl, N₃C₂, and CF₃COO)

COSTES J.-P., DAHAN F., DUPUIS A.

Inorg. Chem., 2000, **39**, pp. 165-168 (LCC-5377).

Is ferromagnetism an intrinsic property of the Cu^{II}/Gd^{III} couple? 1. Structures and magnetic properties of two novel dinuclear complexes with a μ -phenolato- μ -oximate (Cu, Gd) core

COSTES J.-P., DAHAN F., DUPUIS A., LAURENT J.-P.

Inorg. Chem., 2000, **39**, pp. 169-173 (LCC-5378).

Catalysis and kinetics of hydrogen peroxide disproportionation by dinuclear manganese(III) complexes of 1,5-bis(salicylideneamino)pentan-3-ol and the 5-bromophenyl-substituted derivative

PALOPOLI C., CHANSOU B., TUCHAGUES J.-P., SIGNORELLA S.

Inorg. Chem., 2000, **39**, pp. 1458-1462 (LCC-5441).

Synthesis, structure, and magnetic properties of the low-symmetry tetranuclear cubane-like nickel complex $[Ni_4(py-pentO)(pym)(\mu_3-OH)_2(\mu-Oac)_2(NCS)_2(OH_2)]$

CLEMENTE-JUAN J.M., CHANSOU B., DONNADIEU B., TUCHAGUES J.-P.

Inorg. Chem., 2000, **39**, pp. 5515-5519 (LCC-5379).

Is ferromagnetism an intrinsic property of the Cu^{II}/Gd^{III} couple? 2. Structures and magnetic properties of novel trinuclear complexes with μ -phenolato- μ -oximato (Cu-Ln-Cu) cores (Ln = La, Ce, Gd)
COSTES J.-P., DAHAN F., DUPUIS A.

Inorg. Chem., 2000, **39**, pp. 5994-6000 (LCC-5443).

Optical, magnetic and spectroscopic characterization of a novel iron(II) spin-transition compound with a Schiff base ligand

WANG H.M., CHENG P., LI L.C., YAN S.-P., JIANG Z.-H., LIAO D.-Z., WANG G.-L., TUCHAGUES J.-P.

Inorg. Chem. Commun., 2000, **3**, pp. 198-201 (LCC-5444).

Experimental evidence of the acidic character of lanthanide ions in protic solvents

COSTES J.-P., DAHAN F., DUPUIS L., LAURENT J.-P.

Inorg. Chim. Acta, 2000, **298**, pp. 256-259 (LCC-5407).

Infrared detection of the hysteresis in the thermally induced spin-crossover in bis(4,4'-bis-1,2,4-triazole)-bis(thiocyanato-N)iron(II) monohydrate

ZILVERENTANT C.L., VAN ALBADA G.A., BOUSSEKSOU A., HAASNOOT J.G., REEDIJK J.

Inorg. Chim. Acta, 2000, **303**, pp. 287-290 (LCC-5445).

Targeting of a hydrophilic photosensitizer by use of internalizing monoclonal antibodies : a new possibility for use in photodynamic therapy

VROUENRAETS M.B., VISSER G.W.M., LOUP C., MEUNIER B., STIGTER M., OPPELAAR H., STEWART F.A., SNOW G.B., VAN DONGEN G.A.M.S.

Int. J. Cancer, 2000, **88**, pp. 108-114 (LCC-5380).

Guanine oxidation in double-stranded DNA by Mn-TMPyP/KHSO₅: 5,8-dihydroxy-7,8-dihydroguanine residue as a key precursor of imidazolone and parabanic acid derivatives

VIALAS C., CLAPAROLS C., PRATVIEL G., MEUNIER B.

J. Am. Chem. Soc., 2000, **122**, pp. 2157-2167 (LCC-5498).

Rapid synthesis of phosphorus-containing dendrimers with controlled molecular architectures : first example of surface-block, layer-block, and segment-block dendrimers issued from the same dendron

MARAVAL V., LAURENT R., DONNADIEU B., MAUZAC M., CAMINADE A.-M., MAJORAL J.-P.

J. Am. Chem. Soc., 2000, **122**, pp. 2499-2511 (LCC-5381).

A unique coordination of SiH₄ : isolation, characterization, and theoretical study of (PR₃)₂H₂Ru(SiH₄)RuH₂(PR₃)₂

ATHEAUX I., DONNADIEU B., RODRIGUEZ V., SABO-ETIENNE S., CHAUDRET B., HUSSEIN K., BARTHELAT J.-C.

J. Am. Chem. Soc., 2000, **122**, pp. 5664-5665 (LCC-5428).

Ruthenium assisted reversible proton transfer from an aromatic carbon to a hydride

TONER A.J., GRUNDEMANN S., CLOT E., LIMBACH H.-H., DONNADIEU B., SABO-ETIENNE S., CHAUDRET B.

J. Am. Chem. Soc., 2000, **122**, pp. 6777-6778 (LCC-5429).

Experimental evidence of structural evolution in ultrafine cobalt particles stabilized in different polymers - From a polytetrahedral arrangement to the hexagonal structure

DASSENOY F., CASANOVE M.-J., LECANTE P., VERELST M., SNOECK E., MOSSET A., OULD ELY T., AMIENS C., CHAUDRET B.

J. Chem. Phys., 2000, **112**, pp. 8137-8145 (LCC-5193).

Synthesis and X-ray characterization of a monomeric Cp-free d¹-imido-vanadium(IV) complex

LORBER C., DONNADIEU B., CHOUKROUN R.

J. Chem. Soc., Dalton Trans., 2000, pp. 4497-4498 (LCC-5386).

Synthesis, structure and properties of new unsymmetrical nickel dithiolene complexes useful as near-infrared dyes

BIGOLI F., CASSOUX P., DEPLANO P., MERCURI M.L., PELLINGHELLI M.-A., PINTUS G., SERPE A., TROGU E.F.

J. Chem. Soc., Dalton Trans., 2000, pp. 4639-4644 (LCC-5451).

A straightforward preparation of primary alkyl and their utility in the synthesis of derivatives of ethidium

ROSS S.A., PITIE M., MEUNIER B.

J. Chem. Soc., Perkin Trans.1, 2000, pp. 571-574 (LCC-5497).

Alkylating properties of synthetic trioxanes related to artemisinin

CAZELLES J., CAMUZAT-DEDENIS B., PROVOT O., ROBERT A., MAYRARGUE J., MEUNIER B.

J. Chem. Soc., Perkin Trans.1, 2000, pp. 1265-1270 (LCC-5383).

Synthesis and DNA cleavage of 2'-O-amino-linked metalloporphyrin-oligonucleotide conjugates

DUBEY I., PRATVIEL G., MEUNIER B.

J. Chem. Soc., Perkin Trans.1, 2000, pp. 3088-3095 (LCC-5387).

Synthesis and reactivity studies of model complexes for molybdopterin-dependent enzymes

THAPPER A., LORBER C., FRYXELIUS J., BEHRENS A., NORDLANDER E.

J. Inorg. Biochem., 2000, **79**, pp. 67-74 (LCC-5452).

Toward model complexes of Co-containing nitrile hydratases : synthesis, complete characterization and reactivity toward ligands such as CN⁻ and NO of the first square planar Co^{III} complex with two different carboxamido nitrogens and two thiolato sulfur donors

CHATEL S., RAT M., DIJOLS S., LEDUC P., TUCHAGUES J.-P., MANSUY D., ARTAUD I.

J. Inorg. Biochem., 2000, **80**, pp. 239-246 (LCC-5446).

Low-temperature crystal structures of tetrakis- μ -3,5-diisopropylsalicylatobis-dimethylformamidocopper(II) and tetrakis- μ -3,5-diisopropylsalicylatobis-diethyletheratodicopper(II) and their role in modulating polymorphonuclear leukocyte activity in overcoming seizures

MORGANT G., DUNG N.-H., DARAN J.-C., VIOSSAT B., LABOUZE X., ROCH-ARVEILLER M., GREENAWAY F.T., CORDES W., SORENSON J.R.J.

J. Inorg. Biochem., 2000, **81**, pp. 11-22 (LCC-5472).

Synthesis, crystal structure and solid state NLO properties of a new chiral bis(salicyldiaminato)nickel^{II} Schiff-base complex in a nearly optimized solid state environment

AVERSENG F., LACROIX P.G., Malfant I., DAHAN F., NAKATANI K.

J. Mater. Chem., 2000, **10**, pp. 1013-1018 (LCC-5447).

Synthesis and study of radical cation salts and TCNQ charge transfer complexes of a series of tetrathiafulvalenes (TTF) substituted by one or two hydroxylated side chain(s) : -SCH₂CH₂OH

LEGROS J.-P., DAHAN F., BINET L., CARCEL C., FABRE J.-M.

J. Mater. Chem., 2000, **10**, pp. 2685-2691 (LCC-5479).

Convenient route for the preparation of unsymmetrical phospholes via zirconacyclopentadienes

HYDRIO J., GOUYGOU M., DALLEMER F., DARAN J.-C., BALAVOINE G.G.A.

J. Organomet. Chem., 2000, **595**, pp. 261-267 (LCC-5473).

Synthesis, characterization and solution studies on heteronuclear iron-copper clusters containing diphosphine ligands : crystal structure of the dicluster [$\{Fe_2(\mu-CO)(CO)_6(\mu-PPh_2)Cu\}_2(dppp)$]

FERRER M., ROSSELL O., SECO M., SOLER M., FONT-BARDIA M., SOLANS X., DE MONTAUZON D.

J. Organomet. Chem., 2000, **598**, pp. 215-221 (LCC-5384).

Selective P-N bond cleavage in chiral triquinphosphoranes promoted by zirconocene complexes

MARCHI C., IGAU A., MAJORAL J.-P., BUONO G.

J. Organomet. Chem., 2000, **599**, pp. 304-307 (LCC-5522).

Ferrocene and ferrocenyl derivatives in luminescent systems

FERY-FORGUES S., DELAVAUX-NICOT B.

J. Photochem. Photobiol. A, 2000, **132**, pp. 137-159 (LCC-5414).

Nanoscale bimetallic $\text{Co}_x\text{Pt}_{1-x}$ particles dispersed in poly(vinylpyrrolidone): synthesis from organometallic precursors and characterization

OULD ELY T., PAN C., AMIENS C., CHAUDRET B., DASSENOY F., LECANTE P., CASANOVE M.-J., MOSSET A., RESPAUD M., BROTO J.-M.

J. Phys. Chem. B, 2000, **104**, pp. 695-702 (LCC-5423).

Evidence for an original electroactive fluorescent ferrocenyl species and approach to its photophysical behaviour

DELAVAUX-NICOT B., FERY-FORGUES S., MARCOTTE N.

Luminescence, 2000, pp. 59-60 (LCC-5415).

Surface Characterization of Polymer-Stabilized Colloidal Metal Catalysts

DE CARO D., KOHLER J., BUSSE W., BRADLEY J.S.

Macromol. Symp., 2000, **156**, pp. 53-60 (LCC-6664).

Surface characterization of polymer-stabilized colloidal metal catalysts

DE CARO D., KOHLER J., BUSSE W., BRADLEY J.S.

Macromol. Symp., 2000, **156**, pp. 53-60 (LCC-5450).

Phosphorus-containing dendrimers with ferrocenyl units at the core, within the branches, and on the periphery

TURRIN C.O., CHIFFRE J., DE MONTAUZON D., DARAN J.-C., CAMINADE A.-M., MANOURY E., BALAVOINE G.G.A., MAJORAL J.-P.

Macromolecules, 2000, **33**, pp. 7328-7336 (LCC-5388).

Choice of strategies for the divergent synthesis of phosphorus-containing dendrons, depending on the function located at the core

MARAVAL V., PREVOTE-PINET D., LAURENT R., CAMINADE A.-M., MAJORAL J.-P.

New J. Chem., 2000, **24**, pp. 561-566 (LCC-5389).

Synthesis of symmetrical and asymmetrical singly bridged bis-TTFs containing sulfur and selenium atoms. A study of some of their radical cation salts

CARCEL C., FABRE J.-M., GARREAU DE BONNEVAL B., COULON C.

New J. Chem., 2000, **24**, pp. 919-924 (LCC-5449).

Unprecedented amido-bridged dinuclear mixed-valence cobalt complexes : first structure of a dinuclear $\mu\text{-OMe Co(II)/Co(III)}$ complex

HEMMERT C., GORNITZKA H., MEUNIER B.

New J. Chem., 2000, **24**, pp. 949-951 (LCC-5390).

Mechanisms of DNA cleavage by copper complexes of 3Clip-Phen and of its conjugate with a distamycin analogue

PITIE M., BURROWS C.J., MEUNIER B.

Nucleic Acids Res., 2000, **28**, pp. 4856-4864 (LCC-5391).

Five-coordinate organo-zirconocene-ate complexes : synthesis and reactivity

MIQUEL Y., CADIerno V., DONNADIEU B., IGAU A., MAJORAL J.-P.

Organometallics, 2000, **19**, pp. 54-61 (LCC-5519).

Synthesis and characterization of $[\text{Cp}_2\text{V}(\mu\text{-}h^2\text{:}h^4\text{-butadiyne})\text{ZrCp}'_2]$ heterodimetallic complexes ($\text{Cp}' = \text{C}_5\text{H}_4\text{t-Bu}$, $\text{C}_5\text{H}_4\text{Me}$). Formation mechanism and theoretical (ELF) evidence for the existence of planar tetracoordinate carbon (ptC)

CHOUKROUN R., DONNADIEU B., ZHAO J.S., CASSOUX P., LEPETIT C., SILVI B.

Organometallics, 2000, **19**, pp. 1901-1911 (LCC-5392).

Synthesis and structure of group 4 and metal complexes with an ancillary sterically demanding diamido ligand

LORBER C., DONNADIEU B., CHOUKROUN R.
Organometallics, 2000, **19**, pp. 1963-1966 (LCC-5393).

Ruthenium dihydridobis(pyrazolyl)borate complexes adopting a $\kappa^3\text{N,N,H}$, $\kappa^2\text{N,H}$, or $\kappa^2\text{N,N}$ bonding mode

RODRIGUEZ V., ATHEAUX I., DONNADIEU B., SABO-ETIENNE S., CHAUDRET B.
Organometallics, 2000, **19**, pp. 2916-2926 (LCC-5427).

New ferrocenyloxazoline for the preparation of ferrocenes with planar chirality

MANOURY E., FOSSEY J.S., AIT-HADDOU H., DARAN J.-C., BALAVOINE G.G.A.
Organometallics, 2000, **19**, pp. 3736-3739 (LCC-5474).

Highly diastereoselective preparation of ruthenium bis(diimine) sulfoxide complexes : new concept in the preparation of optically active octahedral ruthenium complexes

PEZET F., DARAN J.-C., SASAKI I., AIT-HADDOU H., BALAVOINE G.G.A.
Organometallics, 2000, **19**, pp. 4008-4015 (LCC-5478).

Phosphorus-containing dendrimers and their transition metal complexes as efficient recoverable multicenter homogeneous catalysts in organic synthesis

MARAVAL V., LAURENT R., CAMINADE A.-M., MAJORAL J.-P.
Organometallics, 2000, **19**, pp. 4025-4029 (LCC-5394).

Synthesis and X-ray crystal structure of a urana[1]ferrocenophane, the first tris(1,1'-ferrocenylene) metal compound

BUCAILLE A., LE BORGNE T., EPHRITIKHINE M., DARAN J.-C.
Organometallics, 2000, **19**, pp. 4912-4914 (LCC-5475).

Ruthenium-catalyzed silylation of ethylene by disilanes

DELPECH F., MANSAS J., LEUSER H., SABO-ETIENNE S., CHAUDRET B.
Organometallics, 2000, **19**, pp. 5750-5757 (LCC-5494).

Chlorine free synthesis of organophosphorus compounds based on the functionalization of white phosphorus (P_4)

ABDREIMOVA R.R., AKBAYEVA D.N., POLIMBETOVA G.S., CAMINADE A.-M., MAJORAL J.-P.
Phosphorus Sulfur Silicon Relat. Elem., 2000, **156**, pp. 239-254 (LCC-5395).

Macrocycles à atomes de phosphore et de bore hypervalents. Détermination de leur taille par spectrométrie de masse

RICHELME S., CLAPAROLS C., LEROY E., CAMINADE A.-M., MUNOZ A.
Phosphorus Sulfur Silicon Relat. Elem., 2000, **161**, pp. 143-156 (LCC-5396).

Magnetic properties of $1\text{-BETS}_2(\text{Fe}_2\text{Ga}_{1-x})\text{Cl}_4$ exhibiting a superconductor-to-insulator transition ($0,35_x0,5$)

SATO A., OJIMA E., AKUTSU H., NAKAZAWA Y., KOBAYASHI H., TANAKA H., KOBAYASHI A., CASSOUX P.
Phys. Rev. B-Cond. Matt., 2000, **61**, pp. 111-114 (LCC-5397).

High magnetic field induced spin transition, H.M.F.I.S.T. effect in $[\text{Fe}_{0.52}\text{Ni}_{0.48}(\text{btr})_2(\text{NCS})_2]\text{H}_2\text{O}$

NEGRE N., GOIRAN M., BOUSSEKSOU A., HAASNOOT J.G., BOUKHEDDADEN K., ASKENAZY S., VARRET F.
Synth. Met., 2000, **115**, pp. 289-292 (LCC-5505).

Synthesis of functionalized mono-, tri-, and tetraphosphines: attempted application to prepare hyperbranched polymers and dendrimers built with phosphines at each branching point

HUC V., BALUEVA A., SEBASTIAN R.M., CAMINADE A.-M., MAJORAL J.-P.
Synthesis, 2000, pp. 726-730 (LCC-5398).

Some aspects of the structural and coordination chemistry of the 2,2'-biphosphole
TISSOT O., HYDRIO J., GOUYGOU M., DALLEMER F., DARAN J.-C., BALAVOINE G.G.A.
Tetrahedron, 2000, **56**, pp. 85-93 (LCC-5476).

Dendrimers with N,N-disubstituted hydrazines as end groups, useful precursors for the synthesis of water-soluble dendrimers capped with carbohydrate, carboxylic or boronic acid derivatives
SEBASTIAN R.M., MAGRO G., CAMINADE A.-M., MAJORAL J.-P.
Tetrahedron, 2000, **56**, pp. 6269-6277 (LCC-5399).

Synthesis of chiral molecules containing pyridine and 1,3-pyrimidine units : potential building blocks for helixating and caging ligands
PEZET F., ROUTABOUL L., DARAN J.-C., SASAKI I., AIT-HADDOU H., BALAVOINE G.G.A.
Tetrahedron, 2000, **56**, pp. 8489-8493 (LCC-5477).

Vicinal di-functionalization of α , β -unsaturated ketones via manganese carbene intermediates: synthesis of γ -acyl- δ -lactones by cascade Michael addition/aldol/tranesterification reactions
MONGIN C., GRUET K., LUGAN N., MATHIEU R.
Tetrahedron Lett., 2000, **41**, pp. 7341-7345 (LCC-5419).